



Paper Type: Original Article

Towards an Info-Geometric Theory of the Analysis of Non-Time Dependent Queueing Systems

Ismail A Mageed* 

PhD, AIMMA, IEEE, IAENG, School of Computer Science, AI, and Electronics, Faculty of Engineering and Digital Technologies, University of Bradford, United Kingdom; drismail664@gmail.com.

Citation:

Received: 16 August 2024
Revised: 28 October 2024
Accepted: 20 December 2024

A Mageed, I. (2024). Towards an info-geometric theory of the analysis of non-time dependent queueing systems. *Risk assessment and management decisions*, 1(1), 154-197.


Abstract

Information Geometry (IG) offers a differential geometric perspective for examining statistical models, where families of probability distributions are regarded as manifolds. This paper presents a new method for modeling the IG of a stable M/G/1 queueing system. By defining the M/G/1 queue manifold, the Fisher Information Matrix (FIM) and its inverse (IFIM) are derived, demonstrating the critical influence of system stability on the existence of IFIM. Divergence measures like Kullback's Divergence (KD) and J-Divergence (JD) are explored, revealing the effects of server utilization and the squared coefficient of variation on these measures. It is established that the stable M/G/1 queue manifold is developable (with zero Gaussian curvature) while containing a non-zero Ricci Curvature Tensor (RCT), along with new dynamics that connect RCT behavior to different stability phases. The Information Matrix Exponential (IME) is introduced, illustrating its instability and its reverse correlation with the stability of the M/G/1 manifold. The investigation is broadened to include other divergence measures, merging Queueing Theory, IG, and Riemannian Geometry. Connections to advanced machine learning and metric learning are examined, emphasizing queue learning as an innovative approach. A comprehensive analytical exploration incorporates Gaussian and Ricci curvatures, the Einstein Tensor, and the Stress-Energy Tensor, offering insights into their stability dynamics and geometric significance. For the first time, this research provides thorough derivations of Gaussian curvature, RCT, and their relationships with stability analysis. Computational IG is employed to visualize queueing systems, forging new links between Queueing Theory, matrix theory, IG, and the Theory of Relativity. This cohesive framework enhances both theoretical and practical understandings of queueing systems.

Keywords: Einstein Tensor, Stress Energy Tensor, Riemannian metric, Inverse fisher information matrix, sAB divergence, QT-IG unifiers.

1 | Introduction

Information Geometry (IG) is a field that applies techniques from differential geometry to statistics [1]. It aims to use geometric metrics to provide a new way to describe the probability density function, serving as a coordinate system in Statistical Manifolds (SMs). A manifold [2] is a mathematical concept that represents a space with certain properties. In this context, a manifold is a finite-dimensional Cartesian space, denoted as

 Corresponding Author: drismail664@gmail.com



 Licensee System Analytics. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0>).

\mathbb{R}^n , where \mathbb{R}^n refers to a topological space. It is important to note that although figures can be visualized, they are considered abstract geometric figures rather than concrete representations.

In the given context, IG is highlighted as being significantly important [1], [3], [4]. Fig. 1 illustrates how parameter inference, represented by $\hat{\theta}$. Additionally, previous research has explored the geometric structures of exponential distribution families.




Fig. 1. SM's parametrization [3].

One mathematical method for solving systems of linear differential equations is the Information Matrix Exponential (IME). It also has significant applications in the theory of Lie groups, which are mathematical structures that have important implications in various areas of mathematics and physics [3]. Interarrival time distribution (IG) of stable M/D/1 queues was studied by using features of queue length pathways; the article introduced a geometric structure to the set of M/D/1 queues; for a more detailed survey, consult [3]. This strategy connected information matrix theories with IG, opening new insights into queueing theory. According to [3], [5], Ricci curvature quantifies the distinction between the standard Euclidean Metric (EM) and the Riemannian Metric (RM) in the setting of the article. In contrast, the difference in volume between a geodesic ball and an Euclidean ball with the same radius is measured by scalar curvature. Fig. 2 depicts the knowledge facilitation of comprehension of the geometric characteristics of spaces and how they deviate from Euclidean geometry.




Fig. 2. curved surfaces' geodesic representation [6].

I. states that the exponential of the Fisher Information Matrix (FIM) for the stable M/G/1 queue, a mathematical model used in queueing theory, solves $dx/dt = Ax$. Here, x represents a vector with n dimensions, and A is a $n \times n$ matrix. The second extended study by [6] builds upon their previous work [7] and introduces new contributions:

II. Finding the underlying QM's KD and JD measures.

III. Proving that the FIM of the underlying QM solves: $dx/d = Ax$.

This current paper is an ultimate extension of both papers, with the main deliverables :

I. Extending the study over two new additional divergence measures, namely Rényi's and SAB's, together with a complete illustrative numerical result for all these measures, including KD, JD. This links Queueing theory

- IG with deep machine learning and metric learning. Furthermore, this reveals the revolutionary approach of queue learning.
- II. The solenoidability (incompressibility) of the underlying queueing system is shown. This concept is analogous to the divergence theorem [8].
 - III. Full analytic study of Gaussian curvatures subject to both Angular and Monge Techniques together with the overall stability dynamics impact on these curvatures.
 - IV. The current paper provides a comprehensive analysis of the Einstein Tensor and Stress Energy Tensor, exploring their relationship with stability dynamics and curvatures. It also introduces the definitions and interpretations of Gaussian and Ricci curvatures, as well as the Einstein Tensor.
 - V. Extending the study to include two new divergence measures, Rényi's and *sAB's*, along with illustrative numerical results such as KD and JD. This extension connects queueing theory, IG, deep machine learning, and metric learning, revealing a novel approach called queue learning.
 - VI. Additionally, the paper explores the impact of stability dynamics on Gaussian curvatures, provides a comprehensive analysis of Einsteinian and Stress Energy Tensors, and establishes a unified theorem of queueing-theoretic correlations with both special and general relativity.

The road map of this study is as follows: the core definitions for IG are contained in Section 2. In Section 3, FIM and its inverse for the underlying QM are obtained. In Section 4, the α -connection of a stable M/G/1 queue manifold is obtained. In Section 5, the KD and JD [6], Rényi's, and *sAB* divergences of a stable M/G/1 QM are computed. In Section 6, methodical arguments are developed, demonstrating the developability and non-zero RCT for the underlying queueing manifold system. In addition, a comprehensive examination of the recently announced QT-IG unifiers is presented in Section 6. Section 7 investigates $e^{\text{FIM}(M/G/1)}$ and how the underlying QM's stability impacts FIM's stability.

In Section 8, Ricci scalar, \mathcal{R} , a curvature of spacetime (Einstein tensor) \wp , stress-energy tensor, T , the corresponding threshold theorems for the underlying curvatures, and the dual queueing impact on the existence of the inverse Fisher Information Matrix (IFIM). Section 9 discusses the cueing theoretic impact on the continuity of newly devised Queueing-Information Geometric Unifiers (QIGU). Section 10 is entirely devoted to closing remarks combined with the next phase of research.

2 | Main Definitions

2.1 | Main Definition of IG

Definition 1 ([6]). SM. We denote a SM, $M = \{p(x, \theta) | \theta \in \Theta\}$ and $p(x, \theta)$ as a PDF. Here, $\theta = (\theta_1, \theta_2, \dots, \theta_n) \in \Theta \subset \mathbb{R}^n$.

Definition 2 ([6]). Potential function. The potential function $\Psi(\theta)$ denotes $(-\mathcal{L}(x; \theta) = -\ln(p(x; \theta)))$ with coordinates only.

Definition 3. FIM, namely $[g_{ij}]$. $[g_{ij}]$ [6] reads as

$$[g_{ij}] = \left[\frac{\partial^2}{\partial \theta^i \partial \theta^j} (\Psi(\theta)) \right], i, j = 1, 2, \dots, n. \quad (1)$$

Definition 4. IFIM, namely $[g^{ij}]$ [6].

$$[g^{ij}] = ([g_{ij}])^{-1} = \frac{\text{adj}[g_{ij}]}{\Delta}, \Delta = \det[g_{ij}]. \quad (2)$$

The arc length is defined to be:

$$(ds)^2 = \sum_{i,j=1}^n g_{ij} (d\theta^i)(d\theta^j). \quad (3)$$

Definition 5. α -Connection [6] The α -connection reads as

$$\Gamma_{ij,k}^{(\alpha)} = \left(\frac{1-\alpha}{2}\right)(\partial_i \partial_j \partial_k(\Psi(\theta))), \partial_i = \frac{\partial}{\partial \theta_i}, \alpha \text{ is real.} \tag{4}$$

Definition 6. Kullback's Divergence (KD), $K(p, q)$. KD, namely $K(p, q)$ [6] reads as

$$K(p, q) = E_{\theta_p} \left[\ln \left(\frac{p(x; \theta_p)}{q(x; \theta_q)} \right) \right] = \int p(x; \theta_p) \ln \left(\frac{p(x; \theta_p)}{q(x; \theta_q)} \right) dx. \tag{5}$$

J-Divergence (JD) reads as

$$J(p, q) = \int \ln \left(\frac{p(x; \theta_p)}{q(x; \theta_q)} \right)^{(p(x; \theta_p) - q(x; \theta_q))} dx = K(p, q) + K(q, p). \tag{6}$$

In this paper, however, we focus on the Rényi divergence [9], [10].

$$D_R^\gamma(p||q) = \frac{1}{(\gamma - 1)} \ln \left(\sum_{n=0}^{\infty} (p(n))^\gamma (q(n))^{1-\gamma} \right). \tag{7}$$

used in Rényi variational inference VI [11].

$D_{s,AB}^{\gamma,\eta}(p||q)$ [12] reads as

$$D_{s,AB}^{\gamma,\eta}(p||q) = \frac{1}{\eta(\eta+\gamma)} \ln \left(\sum_{n=0}^{\infty} (p(n))^{\eta+\gamma} \right) + \frac{1}{\gamma(\eta+\gamma)} \ln \left(\sum_{n=0}^{\infty} (q(n))^{\eta+\gamma} \right) - \frac{1}{\gamma\eta} \ln \left(\sum_{n=0}^{\infty} (p(n))^\gamma (q(n))^\eta \right). \tag{8}$$

for $(\gamma, \eta) \in \mathbb{R}^2$ such that $\gamma \neq 0, \eta \neq 0$ and $\gamma + \eta \neq 0$.

The authors [12] have presented a novel (dis)similarity measure, namely $D_{s,AB}^{\gamma,\eta}(p||q)$ Eq. (8). Moreover, it has been illustrated [12] that $D_{s,AB}^{\gamma,\eta}(p||q)$ is potentially robust.




Fig. 3. Beyond euclidean [13].

Definition 7. The α - curvature Riemannian Tensors, $R_{ijkl}^{(\alpha)}$ [6] reads.

$$R_{ijkl}^{(\alpha)} = \left[\left(\partial_j \Gamma_{ik}^{s(\alpha)} - \partial_i \Gamma_{jk}^{s(\alpha)} \right) g_{sl} + \left(\Gamma_{jt,l}^{(\alpha)} \Gamma_{ik}^{t(\alpha)} - \Gamma_{it,l}^{(\alpha)} \Gamma_{jk}^{t(\alpha)} \right) \right], i, j, k, l, s, t = 1, 2, 3, \dots, n, \tag{9}$$

where $\Gamma_{ij}^{k(\alpha)} = \Gamma_{ij,s}^{(\alpha)} g^{sk}, i, j, k, s = 1, 2, \dots, n$.

The α - Ricci curvatures (Ricci Tensors) $R_{ik}^{(\alpha)}$ reads [6].

$$R_{ik}^{(\alpha)} = R_{ijkl}^{(\alpha)} g^{jl}, i, j, k, l = 1, 2, 3, \dots, n. \tag{10}$$

The α – sectional curvatures $K_{ijij}^{(\alpha)}$ reads [6].

$$K_{ijij}^{(\alpha)} = \frac{R_{ijij}^{(\alpha)}}{(g_{ii})(g_{jj}) - (g_{ij})^2}, i, j = 1, 2, \dots, n. \quad (11)$$

Potentially,

$$K^{(\alpha)} = \frac{R_{1212}^{(\alpha)}}{\det(g_{ij})}. \quad (12)$$

One mathematical object that can be obtained by contracting the Riemannian Tensor is the Ricci Tensor [14]. It measures the curvature of space and is employed in the study of Riemannian manifolds. To obtain the Ricci Tensor, the contraction procedure entails summing a few components of the Riemannian Tensor [6].

An oriented Riemannian manifold's Ricci Curvature Tensor (RCT) [15] quantifies the difference between a geodesic ball's volume on the manifold and its volume in Euclidean space. It gives details on the manifold's curvature and how it differs from flat space. Knowledge of the geometry and characteristics of curved spaces in connection to Euclidean geometry requires a knowledge of this topic.

RCT [16] measures how volumes change over time along geodesic paths on a Riemannian manifold. When the Ricci curvature is positive, it indicates a smaller diameter. This relationship is supported by the Bonnet-Myers theorem, which establishes a connection between the manifold's positive Ricci curvature and the curvature properties.




Fig. 4. RCT [17].

Definition 8. Considering the linear system of differential equations.

$$\frac{dx}{dt} = Ax. \quad (13)$$

with x is an n -dimensional vector and A is a $n \times n$ matrix. It can be shown [18] the matrix exponential:

$$e^A = \sum_{i=0}^{\infty} \frac{A^i}{i!} = I + A + \frac{A^2}{2!} + \dots + \frac{A^k}{k!} + \dots. \quad (14)$$

is the solution of Eq. (13).

If the characteristic polynomial of A is defined by

$$\Phi(\delta) = \det(A - \delta I). \quad (15)$$

The eigenvalues of A [18] solve:

$$\Phi(\delta) = (\delta) = 0. \quad (16)$$

such that:

$$Ax = \delta x. \quad (17)$$

e^A reads as:

$$e^A = T e^D T^{-1}. \quad (18)$$

where D is the diagonal matrix of eigenvalues of A , and T is the matrix having the corresponding eigenvectors of A as its columns [18].

Definition 9. Developable surfaces are a special kind of ruled surfaces; they have a Gaussian curvature equal to 0 and can be mapped onto the plane surface without distortion of curves: any curve from such a surface drawn onto the flat plane remains the same [19].




Fig. 5. Three kinds of developable surfaces: a. Tangential (on the left), b. Conical (on the centre), and c. Cylindrical (on the right).

Note that curves in bold are directrix or base curves, and straight lines in bold are directors or generating lines (curves) [19].

2.2 | Gaussian and Mean Curvatures, K_G and H Respectively [20]

Definition 10. (Mong Patch Technique). Let K_1 and K_2 be the principal curvatures of a surface patch $\delta(u, v)$. $K_G(\delta)$ is:

$$K_G = K_1 K_2. \quad (19)$$

and its Mean Curvature is:

$$H = \frac{1}{2}(K_1 + K_2). \quad (20)$$

For a Mong patch $z = f(x, y)$, K_G and H are given by

$$K_G = \frac{LN - M^2}{EG - F^2}. \quad (21)$$

and its Mean Curvature is:

$$H = \frac{1}{2} \left(\frac{LG - 2MF + NE}{EG - F^2} \right). \quad (22)$$

with $E = \left(\frac{\partial f}{\partial x} \right)^2$, $F = \frac{\partial f}{\partial x \partial y}$, $G = \left(\frac{\partial f}{\partial y} \right)^2$, $L = \frac{\partial^2 f}{\partial x^2}$, $M = \frac{\partial^2 f}{\partial x \partial y}$, $N = \frac{\partial^2 f}{\partial y^2}$.

2.2.1 | Classification of surface points




Fig. 6. The elliptic paraboloids $z = x^2 + 2y^2$ (to the left) and $z = x^2 - 2y^2$ (to the right) [20].




Fig. 7. Planar points with quite different shapes [20].

A torus, as shown in Fig. 8.




Fig. 8. Torus [20].




Fig. 9. Catenoid [20].

2.3 | Different Approach to Gaussian and Mean Curvatures (Angular Technique) [21]

A new formulation [21] is introduced for Gaussian Curvature K_G and the Mean Curvature is H is defined as

$$K_G = K_1 K_2. \quad (23)$$

And

$$H = \frac{1}{2}(K_1 + K_2). \quad (24)$$

with K_1 and K_2 as the principal curvatures are determined by

$$K_1 = \frac{B_{11}}{(1+(\frac{\partial f}{\partial x_1})^2)^{\frac{3}{2}}}, K_2 = \frac{B_{22}}{(1+(\frac{\partial f}{\partial x_2})^2)^{\frac{3}{2}}}. \quad (25)$$

where $x_3 = f(x_1, x_2)$ defines the shape of the surface. x'_1 and x'_2 are parallel to the directions of the principal curvature, which are rotated through an angle \mathcal{C} with respect to x_1 and x_2 , and

$$\frac{\partial f}{\partial x'_1} = \cos \mathcal{C} \frac{\partial f}{\partial x_1} - \sin \mathcal{C} \frac{\partial f}{\partial x_2}. \quad (26)$$

$$\frac{\partial f}{\partial x'_2} = \sin \mathcal{C} \frac{\partial f}{\partial x_1} + \cos \mathcal{C} \frac{\partial f}{\partial x_2}. \quad (27)$$

$$B_{11} = \frac{\partial^2 f}{\partial x_1^2} \cos^2 \mathcal{C} - 2 \frac{\partial^2 f}{\partial x_1 \partial x_2} \sin \mathcal{C} \cos \mathcal{C} + \frac{\partial^2 f}{\partial x_2^2} \sin^2 \mathcal{C} = \frac{\partial^2 f}{\partial x_1'^2}. \quad (28)$$

$$B_{22} = \frac{\partial^2 f}{\partial x_1^2} \sin^2 \mathcal{C} + 2 \frac{\partial^2 f}{\partial x_1 \partial x_2} \sin \mathcal{C} \cos \mathcal{C} + \frac{\partial^2 f}{\partial x_2^2} \cos^2 \mathcal{C} = \frac{\partial^2 f}{\partial x_2'^2}. \quad (29)$$

The angle \mathcal{C} through which the coordinate frame is rotated to align the axes with the directions of the principal curvature at each point on the surface is given by

$$\tan 2\mathcal{C} = \frac{-2(\frac{\partial^2 f}{\partial x_1 \partial x_2})}{(\frac{\partial^2 f}{\partial x_1^2} - \frac{\partial^2 f}{\partial x_2^2})}. \quad (30)$$

A contour plot of Gaussian curvature indicates where structures occur on a surface. If both principal curvatures are non-zero, the surface is said to have double curvature [21].




Fig. 10. K_G are slab structures-related by the three regions (a), (b), and (c) [21].

2.4 | Well-Defined Functions and Bijective Functions

Definition 11 ([22]). A function is well-defined if it gives the same result when the representation of the input is changed without changing the value of the input.

Definition 12 ([23]). function f is said to be one-to-one, or injective, if and only if $f(x) = f(y)$ implies $x = y$ for all x and y in the domain of f . A function is said to be an injection if it is one-to-one. Alternative: A function is one-to-one if and only if $f(x) \neq f(y)$, whenever $x \neq y$. This is the contrapositive of the definition.

A function f from A to B is called onto, or surjective, if and only if for every $b \in B$ there is an element $a \in A$ such that $f(a) = b$. Alternative: all co-domain elements are covered.

A function f is called a bijection if it is both one-to-one (injection) and onto (surjection).

Definition 13 ([24]). The solution of the cubic equation.

$$a^*w^3 + b^*w^2 + c^*w + d^* = 0. \quad (31)$$

is characterized arbitrarily by

$$y = z - \frac{\varepsilon_3}{z}. \quad (32)$$

$$w = y - \frac{b^*}{3a^*}. \quad (33)$$

$$z = \sqrt[3]{\left(-\frac{\varepsilon_1}{2}\right) \pm \sqrt{\varepsilon_2}}. \quad (34)$$

$$\varepsilon_1 = \frac{2(b^*)^3}{27} + \frac{d^*}{a^*} - \frac{b^*c^*}{3(a^*)^2}. \quad (35)$$

$$\varepsilon_2 = \frac{(\varepsilon_1)^2}{4} + \frac{(\varepsilon_3)^3}{27}, \quad (36)$$

where ε_3 is given by

$$\varepsilon_3 = -\frac{2(b^*)^2}{3(a^*)^2} + \frac{c^*}{a^*}. \quad (37)$$

ε_2 is called the discriminant of the cubic equation.

Preliminary Theorem (PT) [25]. Let f be a function that is defined and differentiable on an open interval (c,d) .

$$\text{If } f'(x) > 0 \text{ for all } x \in (c, d), \text{ then } f \text{ is increasing on } (c, d). \quad (38)$$

$$\text{If } f'(x) < 0 \text{ for all } x \in (c, d), \text{ then } f \text{ is decreasing on } (c, d). \quad (39)$$

2.5 | Stability Analysis for Ordinary Differential Equations

Equilibria are not always stable. Since stable and unstable equilibria play quite different roles in the dynamics of a system, it is useful to be able to classify equilibrium points based on their stability. Suppose that we have a set of autonomous Ordinary Differential Equations (ODEs) written in vector form [26]:

$$\frac{dx}{dt} = f(x). \tag{40}$$

Suppose that x^* is an equilibrium point. By definition, $f(x^*) = 0$.

Theorem 1 ([26]). An equilibrium point x^* of the differential equation, 1 is stable if all the eigenvalues of J^* , the Jacobian evaluated at x^* , have negative real parts. The equilibrium point is unstable if at least one of the eigenvalues has a positive real part.






Eigenvalues	Fixed point	Flow
complex with positive real parts	unstable focus	
complex with negative real parts	stable focus	
real and positive	unstable node	
real and negative	stable node	
one positive and one negative	saddle point	

Fig. 11. Schematic table of eigenvalues-fixed point-flow [26].

2.6 | Scalar Curvature(Ricci Scalar), \mathcal{R} and Einestein Tensor, \wp

The scalar curvature (Ricci Scalar), \mathcal{R} [15] measures RCT’s contraction.

$$\mathcal{R} = R_{ij}^{(\alpha)} g^{ij}, i, j = 1,2,3, \dots \tag{41}$$

The two-dimensional Ricci Scalar, \mathcal{R} [27] is twice as the Gaussian Curvature K_G ,

$$\mathcal{R} = 2K_G = 2 K_1 K_2. \tag{42}$$

provided that K_1 and K_2 are determined by Eq. (42).

The Ricci scalar \mathcal{R} [15] has a similar meaning to K_G ,

$$\mathcal{R} = \lim_{\epsilon \rightarrow 0} \frac{6n}{\epsilon^2} \left[1 - \frac{A_{\text{curved}}(\epsilon)}{A_{\text{flat}}(\epsilon)} \right]. \tag{43}$$

Ricci scalar completely captures the curvature of the surface.

The equations of motion of a classical theory like General Relativity can be derived directly from a suitable action by using the Euler-Lagrange equations, leading to the well-known Einstein equations [28].

$$G_{ij} = R_{ij}^{(\alpha)} - \frac{\mathcal{R}}{2} g_{ij} = \frac{8\pi\wp\varpi_{ij}}{c^4}, \tag{44}$$

where G_{ij} is the Curvature of Spacetime (Einstein tensor), \wp , $R_{ij}^{(\alpha)}$ defines spacetime – RCT, namely g_{ij} , $\mathcal{R} = R_{ij}^{(\alpha)} g^{ij}, i, j = 1,2,3, \dots$, is the Ricci scalar or scalar curvature, \wp is the universal gravitational constant, c is the speed of light, and ϖ_{ij} are the components of the stress-energy tensor, ϖ , as a descriptor of spacetime-matter-energy distributions.

2.7 | Maxima and Minima of Functions of Two Variables

Suppose that (a_1, b_1) is a critical point of $f(x, y)$ (i.e., $\frac{\partial f(a_1, b_1)}{\partial x} = 0 = \frac{\partial f(a_1, b_1)}{\partial y}$). Let's denote [29]:

$$D = D(a_1, b_1) = f_{xx}(a_1, b_1)f_{yy}(a_1, b_1) - [f_{xy}(a_1, b_1)]^2. \tag{45}$$

This provides the critical point categories:

- I. If $D > 0$ and $f_{xx}(a_1, b_1) > 0$, then there is a relative minimum at (a_1, b_1) .
- II. If $D > 0$ and $f_{xx}(a_1, b_1) < 0$, then there is a relative maximum at (a_1, b_1) .
- III. If $D < 0$, then the point (a_1, b_1) is a saddle point.
- IV. If $D = 0$, then the point (a_1, b_1) may be a relative minimum, relative maximum, or a saddle point. Other techniques would need to be used to classify the critical point.

2.8 | Continuous Functions

Theorem 2. A function f is continuous at x_0 if and only if f is defined on an open interval (r, s) containing x_0 and for each $\epsilon > 0$, there is a $\delta > 0$ such that [30]:

$$|f(x) - f(x_0)| < \epsilon. \tag{46}$$

whenever $|x - x_0| < \delta$.

2.9 | The Maclaren’s Series of $\ln(1 - x)$ for x around zero[28].

$$\ln(1 - x) = - \sum_{n=1}^{\infty} \frac{x^n}{n}. \tag{47}$$

3 | The Fim and Its Inverse for the Stable M/G/1 QM

According to [31], the Maximum Entropy (ME) state probability of the generalized geometric solution of a stable M/G/1 queue (Fig. 12), subject to normalization, Mean Queue Length (MQL), L and server utilization, $\rho (< 1)$ is given by




Fig. 12. Schematic of a stable M/G/1 queue.

$$p(n) = \begin{cases} 1 - \rho, & n = 0, \\ (1 - \rho)g^n, & n \geq 1. \end{cases} \tag{48}$$

where $g = \frac{\rho^2}{(L-\rho)(1-\rho)}$, $x = \frac{L-\rho}{L}$ and $L = \frac{\rho}{2} \left(1 + \frac{1+\rho C_s^2}{1-\rho} \right)$, $\rho = 1 - p(0)$ and $\beta = C_s^2$.

$p(n)$ of Eq. (43) reads as:

$$p(n) = \begin{cases} 1 - \rho, & n = 0, \\ \frac{2\rho \left(\frac{1 + \rho\beta}{1 - \rho} - 1 \right)^{n-1}}{\left(\left(\frac{1 + \rho\beta}{1 - \rho} + 1 \right)^n \right)}, & n > 0, \text{ with } \beta = C_s^2. \end{cases} \tag{49}$$

Theorem 3. The underlying QM satisfies:

I.

$$\text{FIM reads as } [g_{ij}] = \begin{pmatrix} \frac{1}{(1-\rho)^2} & 0 \\ 0 & \frac{-1}{(\beta+1)^2} \end{pmatrix}. \tag{50}$$

$$(ds)^2 = \left(\frac{1}{(1-\rho)^2}\right)(d\rho)^2 - \frac{1}{(\beta+1)^2}(d\beta)^2. \quad (51)$$

II.

$$[g^{ij}] \text{ reads as } [g^{ij}] = \frac{\text{adj}[g_{ij}]}{\Delta} = \begin{pmatrix} (1-\rho)^2 & 0 \\ 0 & -(\beta+1)^2 \end{pmatrix}. \quad (52)$$

Proof:

Case 1. $p(0) = 1 - \rho$. Thus,

$$\mathcal{L}(x; \theta) = \ln(p(x; \theta)) = \ln(1 - \rho),$$

$$\theta = \theta_1 = \rho, \quad (53)$$

$$\Psi(\theta) = -\ln(1 - \rho), \quad (54)$$

$$\text{Therefore,} \quad (55)$$

$$\partial_1 = \frac{\partial \Psi}{\partial \rho} = \frac{1}{1-\rho},$$

$$\partial_1 \partial_1 = \frac{\partial^2 \Psi}{\partial \rho^2} = \frac{1}{(1-\rho)^2}. \quad (56)$$

FIM is given by

$$[g_{ij}] = \left[\frac{\partial^2 \Psi}{\partial \rho^2} \right] = \left[\frac{1}{(1-\rho)^2} \right]. \quad (57)$$

$[g^{ij}]$ reads as

$$[g^{ij}] = [g_{ij}]^{-1} = [(1-\rho)^2]. \quad (58)$$

Case 2. For $n > 0$, $p(n) = \frac{2\rho\left(\frac{1+\rho\beta}{1-\rho}-1\right)^{n-1}}{\left(\frac{1+\rho\beta}{1-\rho}+1\right)^n}$. Therefore, the coordinate system is two-dimensional satisfying:

$$\mathcal{L}(x; \theta) = \ln(p(x; \theta)) = \ln(1 - \rho) + \ln 2 - \ln(\beta + 1) + n \ln\left(\frac{\rho(1 + \beta)}{2 + \rho(\beta - 1)}\right), \quad (59)$$

where

$$\theta = (\theta_1, \theta_2) = (\rho, \beta), \quad \text{with } \beta = C_s^2. \quad (60)$$

We have

$$\Psi(\theta) = \ln(\beta + 1) - \ln(1 - \rho) - \ln 2. \quad (61)$$

$$\text{Thus, we have } \partial_1 = \frac{1}{1-\rho}, \partial_2 = \frac{1}{\beta+1}, \partial_{11} = \frac{1}{(1-\rho)^2}, \partial_1 \partial_2 = \partial_2 \partial_1 = 0, \partial_{22} = -\frac{1}{(\beta+1)^2}. \quad (62)$$

$$\text{FIM is given by } [g_{ij}] = \begin{pmatrix} \frac{1}{(1-\rho)^2} & 0 \\ 0 & \frac{-1}{(\beta+1)^2} \end{pmatrix}. \quad (63)$$

This completes the proof of this part.

It could be verified that:

$$(ds)^2 = \left(\frac{1}{(1-\rho)^2}\right)(d\rho)^2 - \frac{1}{(\beta+1)^2}(d\beta)^2. \quad (51)$$

Finally, after some manipulation, it could be shown that:

$$[g^{ij}] = \frac{\text{adj}[g_{ij}]}{\Delta} = \begin{pmatrix} (1-\rho)^2 & 0 \\ 0 & -(\beta+1)^2 \end{pmatrix}. \quad (52)$$

4 | The $\alpha(\text{OR } \nabla^{(\alpha)})\text{-Connection of the M/G/1 QM}$

Eq. 61 implies:

$$\Gamma_{11,1}^{(\alpha)} = \left(\frac{1-\alpha}{(1-\rho)^3}\right), \Gamma_{11,1}^{(\alpha)} = \frac{(1-\alpha)}{(1-\rho)^3}, \quad (64)$$

$$\Gamma_{11}^{1(\alpha)} = \frac{1-\alpha}{(1-\rho)}, \Gamma_{11}^{1(0)} = \frac{1}{(1-\rho)}, \quad (65)$$

$$\Gamma_{22}^{2(\alpha)} = -\frac{1-\alpha}{(1+\beta)}, \quad \Gamma_{22}^{2(0)} = -\frac{1}{(1+\beta)}. \quad (66)$$

Engaging the same logic, RCT's remaining components can be determined.

5 | Variational Inference, KD, JD, Rényi And s AB Divergences of Stable M/G/1 QM

5.1 | KD and JD Divergences of Stable M/G/1 QM

The following theorem characterizes both KD and JD of Eqs. (5) and (6), respectively.

Theorem 4. The underlying QM satisfies:

$$K(p, q) = E_{\theta_p} \left[\ln \left(\frac{p(x; \theta_p)}{q(x; \theta_q)} \right) \right] = \ln \left(\left(\frac{1-\rho_p}{1-\rho_q} \right) \left(\frac{1+\beta_q}{1+\beta_p} \right) \left[\frac{\rho_q(2+\rho_q(\beta_q-1))}{\rho_p(2+\rho_p(\beta_p-1))} \left(\frac{1+\beta_q}{1+\beta_p} \right) \right]^{L_p} \right). \quad (67)$$

L_p defines the mean queue length at p .

Also,

$$JD(p, q) = \ln \left[\left(\frac{\rho_q(2+\rho_q(\beta_q-1))}{\rho_p(2+\rho_p(\beta_p-1))} \left(\frac{1+\beta_q}{1+\beta_p} \right) \right)^{L_p - L_q} \right]. \quad (68)$$

where L_p, L_q defines the mean queue length at p and q , respectively.

Proof: to show (I), the case for $n = 0$ is straightforward. For $n > 0$, It could be verified that, using Eq. (49), we have.

After some few mathematical steps, it could be seen that:

$$\ln \left(\frac{p(n)}{q(n)} \right) = \ln \left(\frac{\rho_p}{\rho_q} \right) + \ln \left(\frac{1-\rho_p}{1-\rho_q} \right) + (n-1) \ln \left(\frac{\rho_p(1+\beta_p)}{\rho_q(1+\beta_q)} \right) + n \ln \left(\frac{(2+\rho_q(\beta_q-1))}{(2+\rho_p(\beta_p-1))} \right). \quad (69)$$

By Eq. (70), it follows that KD will be determined by.

$$K(p, q) = \ln \left[\left(\frac{1-\rho_p}{1-\rho_q} \right) \left(\frac{1+\beta_q}{1+\beta_p} \right) \right] + L_p \left(\ln \left(\frac{\rho_q(2+\rho_q(\beta_q-1))}{\rho_p(2+\rho_p(\beta_p-1))} \left(\frac{1+\beta_p}{1+\beta_q} \right) \right) \right) \quad (70)$$

$$(n = 0, 1, 2, \dots, \text{ and } L_p = \sum_{n=0}^{\infty} np(n)) = \ln \left(\frac{1-\rho_p}{1-\rho_q} \right) \left(\frac{1+\beta_q}{1+\beta_p} \right) \left[\frac{\rho_q(2+\rho_q(\beta_q-1))}{\rho_p(2+\rho_p(\beta_p-1))} \left(\frac{1+\beta_q}{1+\beta_p} \right) \right]^{L_p}$$

This completes the proof.

Following some mathematical steps, Eq. (68) could be easily verified.

Clearly, it follows from Eq. (68) that JD is also zero if and only if $p = q$. This presents a novel result that declares the compressibility of the underlying QM if and only if it is stable or when p and q are identical (i.e., $\rho_p = \rho_q, \beta_p = \beta_q$).

It is observed by Eq. (67) that JD is dependent on ρ_q, β_q and the MQL of Pollaczek-Khinchin Formula of a stable M/G/1 QM at p, L_p (which is dependent on ρ_p and β_p). To examine the impact of L_p on JD, the following experiment is introduced.

IMPACT OF MQL_p ON (-KD), Su(q) = 0.5, SCV(q) = 2




Fig. 13. The impact of MQL_p on (-KD).

Fig. 13 depicts that KD is a negative decreasing function in MQL at p, L_p . This justifies the increase of L_p will have a significant impact on the decrease of KD. In other words, the increase of MQL at p , would enforce the distance between p and q to increase in magnitude.

The compressibility of M/G/1 QM could be identified visually by presenting the following numerical experiment:

IMPACT OF MQL_p ON (-JD), Su(q) = 0.5, SCV(q) = 2, MQL_q = 1.25




Fig. 14. The impact of MQL_p on (-JD).

The findings of Fig. 14, show that the increase of MQL at p impacts the stable M/G/1 QM's solenoidability of. It was observed that M/G/1 QM is solenoidal at the steady state phase of the QM. The increase of L_p such that $L_p \neq L_q$, the stable M/G/1 QM is no longer solenoidal. This shows the direct impact of queuing parameters on the visualization of the regions of solenoidability of the stable M/G/1 QM.

Meanwhile, we have




Fig. 15. $SU(p)$ vs KD .

As observed in Fig. 15, KD decreases and vanishes at $\rho_p = 0.5$. by the increase of ρ_p , KD decreases and tends to $-\infty$ when the underlying $M/G/1$ QM approaches instability ($\rho_p = 1$).




Fig. 16. $SU(p)$ vs JD .

As observed by Fig. 16, for $n = 2$, KD increases for permissible values of $SU(p)$ and starts to decrease at $SU(p) =$ as we go along. Then, it decreases when $SU(p) = 0.4$. Afterwards, KD decreases unsoundly, speeding rapidly to $-\infty$. In this physical interpretation, stability has a significant impact on the behaviour of KD . In principle, it is uncovered that the stability of the $M/G/1$ QM has a significant impact on the performance of KD .

5.2 | Rényi Divergence of Stable $M/G/1$ QM

Theorem 5. The underlying queueing system satisfies:

$$RD(p, q) = D_R^\gamma(p||q)$$

$$= \begin{cases} \frac{1}{(\gamma-1)} (\gamma \ln(1-\rho_p) + (1-\gamma) \ln(1-\rho_q)), & n = 0 \\ \ln \left[\left(\frac{\rho_p(1+\beta_q)(1-\rho_p)}{(1-\rho_q)(1+\beta_p)} \right)^{\rho_p} + \frac{1}{(\gamma-1)} \ln \left[\sum_{n=1}^{\infty} p(n) \left(\frac{\rho_p(1+\beta_p)(2+\rho_q(\beta_q-1))}{\rho_q(2+\rho_p(\beta_p-1))(1+\beta_q)} \right)^{n(\gamma-1)} \right] \right], & n > 0. \end{cases} \quad (71)$$

Proof:

$$RD(p, q) = D_R^\gamma(p||q) = \frac{1}{(\gamma-1)} \ln \left(((p(0))^{\gamma} (q(0))^{1-\gamma}) \right) = \frac{1}{(\gamma-1)} \ln \left(((1-\rho_p)^{\gamma} (1-\rho_q)^{1-\gamma}) \right). \quad (72)$$

It could be easily checked that $D_R^\gamma(p||q)$ of Eq. (72) that:

$$D_R^\gamma(p||q) = \frac{1}{(\gamma-1)} (\gamma \ln(1-\rho_p) + (1-\gamma) \ln(1-\rho_q)). \quad (72.1)$$

as required.

It could be verified that for $n > 0$,

$$RD(p, q) = \ln \left(\left(\frac{(1-\rho_p)(1+\beta_q)}{(1-\rho_q)(1+\beta_p)} \right)^{\rho_p} + \frac{1}{(\gamma-1)} \ln \left(\sum_{n=1}^{\infty} \left(\frac{\rho_p(1+\beta_p)(2+\rho_q(\beta_q-1))}{\rho_q(2+\rho_p(\beta_p-1))(1+\beta_q)} \right)^{n(\gamma-1)} p(n) \right) \right).$$

We are done.

As $\gamma \rightarrow 1$,

$$\lim_{\gamma \rightarrow 1} D_R^\gamma(p||q) = \lim_{\gamma \rightarrow 1} \frac{1}{(\sum_{n=0}^{\infty} (p(n))^{\gamma} (q(n))^{1-\gamma})} \left(\sum_{n=0}^{\infty} [\gamma (p(n))^{\gamma-1} (q(n))^{1-\gamma} + (1-\gamma) (q(n))^{-\gamma} (p(n))^{\gamma}] \right) = \lim_{\gamma \rightarrow 1} \frac{1}{(\sum_{n=0}^{\infty} (p(n))^{\gamma} (q(n))^{1-\gamma})} \sum_{n=0}^{\infty} (1) = \infty.$$

Data Rényi Divergence, RD.

Case 1.




Fig. 17. Dual impact of ITP and SU(p) on (-RD), ITP = SU(q) = 0.5, n = 0.

It is shown by Fig. 17 that RD is drastically decreasing until SU(p) is greater than 1, RD becomes an imaginary number, i.e., the instability of M/G/1 QM occurs, RD becomes imaginary!!!

It is observed that for $n = 0, \gamma \rightarrow 1$ (shannonian phase), $D_R^1(p||q) = \ln \left(\frac{1-\rho_p}{1-\rho_q} \right)$.

Case 2.




Fig. 18. Dual impact of ITP and SU(p) on (RD).

Following Fig. 18, The decreasability of RD is clear because of the dual impact of Su (p) and ITP on RD.

Case 2, RD, n = 2




Fig. 19. More on SU(p) vs RD.

As observed from Fig. 19, RD increases rapidly with the increase of ρ_p . It is expected that RD approaches infinity as ρ_p approaches unity (i.e, M/G/1 QM is unstable at p). To show this, we can take the limit of Eq. (72.1) as ρ_p approaches unity. This directly implies that:

$$D_R^{0.5}(p||\rho_q = 0.5, \beta_q = 3) \rightarrow \ln 0 + 3\ln(3) - 2\ln(p(1)(1 - p(1))). \tag{73}$$

The devised corresponding absolute limiting value of RD is.

$$|D_R^{0.5}(p||\rho_q = 0.5, \beta_q = 3)| \rightarrow |\ln 0 + 3\ln(3) - 2\ln(p(1)(1 - p(1)))| = \infty. \tag{74}$$

5.2 | sAB Divergence, $D_{s,AB}^{\eta,\gamma}(p || q)$ of Stable M/G/1 QM

Before going into details, we need to prove the following important lemma as it is needed in the proofs.

Lemma 1. $p(n)$ of Eq. (49) rewrites to

$$p(n) = \begin{cases} 1 - \rho, & n = 0 \\ \frac{2\rho(1-\rho)}{(1+\beta)} \left(\frac{\rho^2(1+\beta)}{(1-\rho)L} \right)^n, & n > 0, \text{ with } \beta = C_s^2. \end{cases} \tag{75}$$

where $L = \frac{\rho}{2} \left(1 + \frac{1+\rho C_s^2}{1-\rho} \right)$, $\rho = 1 - p(0)$ and $\beta = C_s^2$.

Proof: For $n = 0$, it is immediate by Eq. (49).

As for $n > 0$. By the MQL formula, we have $L = \frac{\rho}{2} \left(1 + \frac{1+\rho\beta}{1-\rho} \right)$. Hence, $(2 + \rho(\beta - 1)) = \frac{2(1-\rho)L}{\rho}$. This implies:

$$\frac{2\rho \left(\frac{1+\rho\beta-1}{1-\rho} \right)^{n-1}}{\left(\frac{1+\rho\beta+1}{1-\rho} \right)^n} = \frac{2\rho \left(\frac{\rho(1+\beta)}{1-\rho} \right)^{n-1}}{\left(\frac{2+\rho(\beta-1)}{1-\rho} \right)^n} = \frac{2\rho \left(\frac{\rho(1+\beta)}{1-\rho} \right)^{n-1}}{\left(\frac{2(1-\rho)L}{\rho} \right)^n} = \frac{2\rho(1-\rho)}{(1+\beta)} \left(\frac{\rho^2(1+\beta)}{(1-\rho)L} \right)^n. \tag{76}$$

By Eq. (76) and Eq. (49), the proof follows.

Theorem 6. $sAB_{n=0}$ divergence vanishes,

$$D_{s,AB}^{\eta,\gamma}(p||q) = 0. \tag{77}$$

Proof: For $n = 0$, $p(n) = 1 - \rho_p$, $q(n) = 1 - \rho_q$. Hence, it follows by Eq. (8) that:

$$D_{s,AB}^{\eta,\gamma}(p||q) = 0 \tag{78}$$

Theorem 7. For $n \neq 0$, sAB divergence is determined by

$$D_{s,AB}^{\eta,\gamma}(p||q) = \ln \left[\left(\sum_{n=1}^{\infty} \left(\frac{\rho_p^2(1+\beta_p)}{(1-\rho_p)L_p} \right)^{n(\gamma+\eta)} \right)^{\frac{1}{\eta(\eta+\gamma)}} \left(\sum_{n=1}^{\infty} \left(\frac{\rho_q^2(1+\beta_q)}{(1-\rho_q)L_q} \right)^{n(\gamma+\eta)} \right)^{\frac{1}{\gamma(\eta+\gamma)}} \left(\sum_{n=1}^{\infty} \left(\frac{\rho_p^2(1+\beta_p)}{(1-\rho_p)L_p} \right)^{n\gamma} \left(\frac{\rho_q^2(1+\beta_q)}{(1-\rho_q)L_q} \right)^{n\eta} \right)^{-\frac{1}{\gamma\eta}} \right]. \tag{79}$$

for $(\gamma, \eta) \in \mathbb{R}^2$ such that $\gamma \neq 0, \eta \neq 0$ and $\gamma + \eta \neq 0$.

Proof: we have

$$p(n) = \frac{2\rho_p(1-\rho_p)}{(1+\beta_p)} \left(\frac{\rho_p^2(1+\beta_p)}{(1-\rho_p)L_p} \right)^n, \quad q(n) = \frac{2\rho_q(1-\rho_q)}{(1+\beta_q)} \left(\frac{\rho_q^2(1+\beta_q)}{(1-\rho_q)L_q} \right)^n.$$

Following Eq. (8),

$$D_{s,AB}^{\eta,\gamma}(p||q) = \left[\frac{1}{\eta(\eta+\gamma)} \ln \left(\sum_{n=1}^{\infty} \left(\frac{2\rho_p(1-\rho_p)}{(1+\beta_p)} \left(\frac{\rho_p^2(1+\beta_p)}{(1-\rho_p)L_p} \right)^n \right)^{\gamma+\eta} \right) + \frac{1}{\gamma(\eta+\gamma)} \ln \left(\sum_{n=1}^{\infty} \left(\frac{2\rho_q(1-\rho_q)}{(1+\beta_q)} \left(\frac{\rho_q^2(1+\beta_q)}{(1-\rho_q)L_q} \right)^n \right)^{\gamma+\eta} \right) - \frac{1}{\gamma\eta} \ln \left(\sum_{n=1}^{\infty} \left(\frac{2\rho_p(1-\rho_p)}{(1+\beta_p)} \left(\frac{\rho_p^2(1+\beta_p)}{(1-\rho_p)L_p} \right)^n \right)^{\gamma} \left(\frac{2\rho_q(1-\rho_q)}{(1+\beta_q)} \left(\frac{\rho_q^2(1+\beta_q)}{(1-\rho_q)L_q} \right)^n \right)^{\eta} \right) \right]. \tag{80}$$

It could be checked that the RHS of Eq. (80) reduces after some lengthy computation to:

$$D_{s,AB}^{\gamma,\eta}(p||q) = \ln \left[\left(\sum_{n=1}^{\infty} \left(\frac{\rho_p^2(1+\beta_p)}{(1-\rho_p)L_p} \right)^{n(\gamma+\eta)} \right)^{\frac{1}{\eta(\eta+\gamma)}} \left(\sum_{n=1}^{\infty} \left(\frac{\rho_q^2(1+\beta_q)}{(1-\rho_q)L_q} \right)^{n(\gamma+\eta)} \right)^{\frac{1}{\gamma(\eta+\gamma)}} \right. \\ \left. \left(\sum_{n=1}^{\infty} \left(\frac{\rho_p^2(1+\beta_p)}{(1-\rho_p)L_p} \right)^{n\gamma} \left(\frac{\rho_q^2(1+\beta_q)}{(1-\rho_q)L_q} \right)^{n\eta} \right)^{-\frac{1}{\gamma\eta}} \right]. \tag{79}$$

Numerical experiment for $D_{s,AB}^{\gamma,\eta}(p||q)$.

Data 1

Following Eq. (79), it can be verified, after some manipulation, that for $\rho_q = 0.5, \beta_p = \beta_q = 2, n = 2$ DITP = $(\gamma, \eta) = (1, 1)$,

$$D_{s,AB}^{1,1}(p||q) = \ln \left[\frac{\sqrt{\frac{\left(\frac{6\rho_p}{2+\rho_p}\right)^2 + \left(\frac{6\rho_p}{2+\rho_p}\right)^4}{\left(\frac{12}{5}\right)^2 + \left(\frac{12}{5}\right)^4}}}{\left(\frac{72\rho_p}{5(2+\rho_p)}\right) + \left(\frac{72\rho_p}{5(2+\rho_p)}\right)^2} \right].$$




Fig. 20. SU(p) vs GABD.

The increasability of the Generalized Alpha Beta Divergence (GABD) as SU(p) increases is obvious from Fig. 20.

Data 2

After some lengthy computation, it could be verified that for the non-extensive information theoretic dual, DITP = $(\gamma, \eta) = (0.5, 0.5), n = 2$.

$$(D_{s,AB}^{(0.5,0.5)}(p||q))_{(\rho_q=0.5, \beta_q=3, \beta_p=2)} = \ln \left(\frac{3 \sum_{n=1}^2 \left(\frac{6\rho_p}{2+\rho_p}\right)^n}{2\rho_p(1-\rho_p) \left(\sum_{n=1}^2 \left(\frac{6\rho_p}{2+\rho_p}\right)^n\right)^4} \right).$$

As $\rho_p \rightarrow 1$ (instability phase of M/G/1 QM), $D_{AB}^{(0.5,0.5)}(p||q_{\rho_q=0.5, \beta_q=0.5}) \rightarrow \infty$.

6 | Investigations of the Developability of THE Stable M/G/1 QM, RICCI Curvature (RCT) tENSOR and QT-IG Unifiers

6.1 | Investigation of Developability of M/G/1 QM and Finding its Ricci Curvature Tensor

Theorem 8. The stable M/G/1 QM.

- I. Has a zero 0-Gaussian curvature, for which the stable M/G/1 QM would be developable.
- II. Has a non-zero Ricci Tensor.
- III. Is the non-developable minimal surface under Monge Technique, with a zero mean curvature.
- IV. Is developable under Angular Technique if and only if M/G/1 QM unstable.
- V. If the underlying QM is unstable, then the Mean Curvature is negative under angular technique. The converse statement is not always true.
- VI. The first principal curvature, K_1 under the Angular Technique satisfies the inequality

$$K_1 < 1. \tag{81}$$
- VII. The second principal curvature, K_2 is β – dependent and is negative under the Angular Technique.
- VIII. Under the Angular Technique, the second principal curvature, K_2 tends to zero as $\beta \rightarrow \infty$.

Proof:

- I. For we must prove:

$$K^{(\alpha)} = \frac{R_{1212}^{(\alpha)}}{\det(g_{ij})} = 0. \tag{82}$$

It could be verified that, $R_{1212}^{(\alpha)} = 0.$ (83)

$$\det(g_{ij}) = -\frac{1}{(\beta+1)^2(1-\rho)^2} \neq 0. \text{ Hence,}$$

$$K^{(\alpha)} = \frac{R_{1212}^{(\alpha)}}{\det(g_{ij})} = 0, \text{ which proves that the underlying QM is developable subject to } \alpha \text{ – Gaussian curvature.}$$

- II. We must prove that:

$$R_{ik}^{(\alpha)} = R_{ijkl}^{(\alpha)} g^{jl} \text{ is non-zero.}$$

We have

$$R_{1212}^{(\alpha)} g^{11} + R_{1112}^{(\alpha)} g^{12} + R_{1211}^{(\alpha)} g^{21} + R_{1212}^{(\alpha)} g^{22}. \tag{84}$$

Engaging the same procedure as in Eq. (82), we have

$$R_{11}^{(\alpha)} = R_{12}^{(\alpha)} = R_{22}^{(\alpha)} = 0(6.3). \quad R_{21}^{(\alpha)} = -\frac{1-\alpha}{(1-\rho)^2}, \quad R_{21}^{(0)} = -\frac{1}{(1-\rho)^2}. \tag{85}$$

Hence, $R_{21}^{(\alpha)} \neq 0.$ The corresponding RCT is given by

$$(RCT) = \begin{pmatrix} 0 & -\frac{1}{(1-\rho)^2} \\ 0 & 0 \end{pmatrix}. \tag{86}$$

As $\rho \rightarrow 1$, $R_{21}^{(0)} \rightarrow -\infty.$ This highlights the significant influence of instability in a specific type of the underlying QM by providing supporting evidence on how RCT is significantly impacted by the stability

analysis of the system. The also shows that ρ , represented by $R_{21}^{(0)}$, affects the behaviour of RCT, and Fig. 21 demonstrates that the stability phase of the M/G/1 QM causes RCT to decrease as ρ increases.




Fig. 21. How RCT is impacted by the stability of M/G/1 queue.

Whereas in Fig. 22, RCT increasability in ρ is caused by the underlying QM's instability.




Fig. 22. How RCT is impacted by the instability of M/G/1 queue.

III. Following Eq. (61) and Eq. (62), it is clear that.

Thus, we have $\partial_1 = \frac{1}{1-\rho} \partial_2 = \frac{1}{\beta+1}$, $\partial_{11} = \frac{1}{(1-\rho)^2}$, $\partial_1 \partial_2 = \partial_2 \partial_1 = 0$, $\partial_{22} = -\frac{1}{(\beta+1)^2}$.

The gaussian curvature

$$K_G = \frac{LN - M^2}{EG - F^2},$$

$$E = (\partial_1)^2 = \left(\frac{1}{1-\rho}\right)^2, F = \partial_1 \partial_2 = 0, G = (\partial_2)^2 = \frac{1}{(\beta+1)^2},$$

$$L = \partial_{11} = \frac{1}{(1-\rho)^2}, M = \partial_{12} = 0, N = \partial_{22} = -\frac{1}{(\beta+1)^2}.$$

And the mean curvature is

$$H = \frac{1}{2} \left(\frac{LG-2MF+NE}{EG-F^2} \right).$$

Therefore, it is obtained that

$$K_G = \frac{LN-M^2}{EG-F^2} = -\frac{\frac{1}{(1-\rho)^2(\beta+1)^2}}{\frac{1}{(1-\rho)^2(\beta+1)^2}} = -1, \rho \neq 1 \text{ (the underlying QM is stable), } \beta \neq -1. \quad (87)$$

($\beta = C_s^2$ is never negative by default).

Since $K_G = -1$, follows by the non-developability of the underlying QM and its minimal surface under the Monge Technique.

The Mean Curvature is

$$H = \frac{1}{2} \left(\frac{LG-2MF+NE}{EG-F^2} \right) = \frac{\frac{1}{2} \left(\frac{1}{(1-\rho)^2(\beta+1)^2} - \frac{1}{(1-\rho)^2(\beta+1)^2} \right)}{\frac{1}{(1-\rho)^2(\beta+1)^2}} = 0. \quad (88)$$

Hence, III is done.

IV. Following the Angular Technique, it can be verified that the calculations of the principal curvatures K_1 and K_2 are determined by

$$K_1 = \frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}}, K_2 = \frac{-(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}}. \quad (89)$$

$$K_G = K_1 K_2 = \frac{-(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}} \frac{(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}}, \quad (90)$$

and

$$H = \frac{1}{2} (K_1 + K_2) = \frac{1}{2} \left(\frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}} - \frac{(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}} \right). \quad (91)$$

The axial rotator angle \mathcal{C} reads as

$$\tan 2\mathcal{C} = \frac{-2(\partial_1 \partial_2)}{(\partial_{11} - \partial_{22})} = 0. \quad (92)$$

This implies, $\mathcal{C} = 0, 2\pi, 4\pi, \dots$

It appears from Eq. (90), that $K_G = 0$ (equivalently, the underlying QM is developable) if and only if

$$\frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}} \frac{(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}} = 0. \quad (93)$$

This implies that:

$$\text{either } (1-\rho) = 0 \text{ (equivalently, } \rho = 1) \text{ or } (1+\beta) = 0. \quad (94)$$

The second possibility ($(1+\beta) = 0$) generates a contradiction as β is never negative.

Moreover,

$$\lim_{\beta \rightarrow \infty} K_G = -\frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}} \lim_{\beta \rightarrow \infty} \frac{(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}} = 0. \quad (95)$$

Linking the findings of Eq. (93) and Eq. (95) completes the proof of IV.

As for V), it has been obtained that

$$H = \frac{1}{2}(K_1 + K_2) = \frac{1}{2} \left(\frac{(1-\rho)}{(1+((1-\rho)^2)^{\frac{3}{2}}} - \frac{(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}} \right). \quad (96)$$

This directly implies

$$H < \frac{1}{2} \frac{(1-\rho)}{(1+((1-\rho)^2)^{\frac{3}{2}}}. \quad (97)$$

It is clear that $1 + ((1-\rho)^2) > 0$ holds for all the possible values of ρ . Consequently, if $(1-\rho) = 0$ or $\rho = 1$ (equivalently, the underlying QM is unstable), it follows that $H < 0$

To prove the necessity condition, assume that $H < 0$. This generates two possibilities:

The first possibility, $\frac{1}{2} \frac{(1-\rho)}{(1+((1-\rho)^2)^{\frac{3}{2}}} < 0$. This implies $\rho > 1$. Hence, M/G/1 QM is unstable.

The second possibility, $\frac{1}{2} \frac{(1-\rho)}{(1+((1-\rho)^2)^{\frac{3}{2}}} > 0$. This implies $\rho < 1$. Hence, M/G/1 QM is stable.

This justifies that the converse statement is not always true. The proof of V) is complete.

To show VI), we have

$$K_1 = \frac{(1-\rho)}{(1+((1-\rho)^2)^{\frac{3}{2}}}. \quad (89)$$

Since the underlying M/G/1 QM is assumed to be stable. Hence, $\rho \in (0,1)$. Thus, we have

$$1 > (1-\rho)^2 > 0 \text{ or } 2 > (1-\rho)^2 + 1 > 1. \text{ Therefore, } 2^{\frac{3}{2}} > (1+((1-\rho)^2)^{\frac{3}{2}} > 1. \text{ Consequently,} \\ \frac{1}{2^{\frac{3}{2}}} < \frac{1}{(1+((1-\rho)^2)^{\frac{3}{2}}} < 1. \text{ This implies } \frac{(1-\rho)}{2^{\frac{3}{2}}} < \frac{(1-\rho)}{(1+((1-\rho)^2)^{\frac{3}{2}}} < (1-\rho) < 1. \quad (98)$$

By Eq. (98), it holds that $K_1 < 1$.

VII. we have by Eq. (89), $K_2 = \frac{-(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}}$, which is, of course, a β – dependent function. The stability of M/G/1 QM enforces the condition $\beta > 1$ to hold. The negativity of K_2 is clear.

VIII. Immediate from Eq. (95).

6.2 | Revealing Novel QT-IG Unifiers and Discovering Their Algebraic Structures

Throughout this section, the following novel unifiers between both queueing theoretic and information geometric structures of the stable M/G/1 QM are established by the following two unifiers,

$$\varphi_1(\rho) = K_1 = \frac{(1-\rho)}{(1+((1-\rho)^2)^{\frac{3}{2}}}. \quad (100)$$

$$\varphi_2(\beta) = K_2 = \frac{-(1+\beta)}{(1+((1+\beta)^2)^{\frac{3}{2}}}. \quad (101)$$

Theorem 9. For the above-devised unifiers Eqs. (100) and (101), it holds that

φ_1 is a well-defined function.

$$\varphi_1 = \begin{cases} \text{lies in the interval } (0,1), & \rho \in (0,1), \\ 0, & \rho = 1, \\ < 0, & \rho > 1. \end{cases} \quad (102)$$

φ_1 is one-to-one.

φ_1 is onto.

φ_1 is a bijection with an imaginary inverse φ_1^{-1} determined by

$$\varphi_1^{-1}(\rho) = 1 \mp \left((z-1) + \frac{(3 + \frac{1}{\rho^2})}{z} \right)^{\frac{1}{2}}. \quad (103)$$

$$\text{where } z = \sqrt[3]{-\frac{1}{2\rho^2} \pm i \sqrt{\frac{1}{12\rho^4} + 1 + \frac{1}{2\rho^2} + \frac{1}{27\rho^6}}}, i = \sqrt{-1}. \quad (104)$$

φ_2 is a well-defined function.

$$\varphi_2 = \begin{cases} < -\frac{2}{3}, & \beta \in (0,1), \\ -\frac{2}{3}, & \beta = 1, \\ > -\frac{2}{3}, & \beta > 1. \end{cases} \quad (105)$$

φ_2 is one-to-one.

φ_2 is onto.

φ_2 is a bijection with an imaginary inverse φ_2^{-1} (i.e., a complex number) determined by

$$\varphi_2^{-1}(\beta) = -1 \mp \left((z-1) + \frac{(3 + \frac{1}{\beta^2})}{z} \right)^{\frac{1}{2}}, \quad (106)$$

$$\text{where } z = \sqrt[3]{-\frac{1}{2\beta^2} \pm i \sqrt{\frac{1}{12\beta^4} + 1 + \frac{1}{2\beta^2} + \frac{1}{27\beta^6}}}, i = \sqrt{-1}, \quad (107)$$

The underlying QM has the inverse of ρ unifier, namely, φ_1^{-1} satisfies.

$$|1 - \varphi_1^{-1}(\rho)| < 5 + \frac{(1 + (\frac{101}{54})^{\frac{1}{6}})}{\rho^2} \quad (108)$$

The underlying of the inverse of β unifier, namely, φ_2^{-1} satisfies.

$$|1 - \varphi_2^{-1}(\beta)| < (1 + (\frac{101}{54})^{\frac{1}{6}}) + \frac{1}{(\frac{101}{54})^{\frac{1}{6}}} + \frac{3\beta^2}{(\frac{101}{54})^{\frac{1}{6}}}. \quad (109)$$

XIV) The increasability and decreasability of φ_1 in ρ are undecidable.

XV) φ_2 is forever increasing in β and is never decreasing in β .

To prove I), it is enough to show that for all $\rho_1, \rho_2 \in (0,1)$ such that $\varphi_1(\rho_1) = \varphi_1(\rho_2)$, then ρ_1 and ρ_2 should never be distinct.

Let $\varphi_1(\rho_1) = \varphi_1(\rho_2)$. After some lengthy mathematical steps, Eq. (100) reduces to:

$$(\rho_1 - \rho_2)(\rho_1 + \rho_2 - 2)[((1 - \rho_1)^2(1 - \rho_2)^2((1 - \rho_1)^2 + (1 - \rho_2)^2 - 2))] = 0. \quad (110)$$

Eq. (110) generates three possible cases:

Case 1. $\rho_1 - \rho_2 = 0$. Hence $\rho_1 = \rho_2$ (contradiction to $\rho_1 \neq \rho_2$). (111)

Case 2. $\rho_1 + \rho_2 - 2 = 0$. Hence $\rho_1 + \rho_2 = 2$ (contradiction, since M/G/1 is a stable QM, $\rho_1, \rho_2 \in (0,1)$). (112)

Case 3. $[((1 - \rho_1)^2(1 - \rho_2)^2((1 - \rho_1)^2 + (1 - \rho_2)^2 - 2))] = 0$. (113)

By $\rho_1, \rho_2 \in (0,1)$, following mathematical analysis, it holds that.

Therefore, $[((1 - \rho_1)^2(1 - \rho_2)^2((1 - \rho_1)^2 + (1 - \rho_2)^2 - 2))] < 0$, which contradicts.

Based on the above analysis and by Eq. (111),

I. follows.

II. we have by Eq. (100), $\varphi_1(\rho) = K_1 = \frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}}$. Since $\rho \in (0,1)$, it could be verified that.

$$2 > 1 + ((1 - \rho))^2 > 0. \text{ Hence, } \frac{1}{(1+((1-\rho))^2)^{\frac{3}{2}}} \in (0, \frac{1}{2}). \text{ Therefore, } 0 < \varphi_1(\rho) = \frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}} < \frac{1}{2} < 1. \quad (114)$$

The case $\varphi_1(1) = 0$ is clear. Also, for $\rho > 1$, it is immediate that $\varphi_1(\rho) < 0$. This completes the proof of II).

III. It suffices to show that for all $\rho_1, \rho_2 \in (0,1)$ such that $\varphi_1(\rho_1) = \varphi_1(\rho_2)$, then $\rho_1 = \rho_2$ holds. The proof is clearly immediate from Eq. (111).

IV. From the definition, $\varphi_1(\rho) = \frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}}$. Every $\frac{(1-\rho)}{(1+((1-\rho))^2)^{\frac{3}{2}}}$ is characterized by ρ . This clearly proves the surjectivity of φ_1 . Hence, IV) follows.

V. Clearly, φ_1 is a bijection. To calculate the inverse of φ_1 , namely φ_1^{-1} . Define $\varphi_1(\rho) = y$. Hence, $\frac{(1+((1-\rho))^2)^3}{y^2} = (1 - \rho)^2$. Let $w = (1 - \rho)^2$. Then, we have the cubic equation:

$$w^3 + 3w^2 + \left(3 - \frac{1}{y^2}\right)w + 1 = 0. \quad (115)$$

Following the method for solving cubic equations, we have

$$a^* = 1, b^* = 3, c^* = \left(3 - \frac{1}{y^2}\right), d^* = 1. \quad (116)$$

The solution of Eq. (115) is characterized arbitrarily by

$$w = r - 1, \quad (117)$$

$$\gamma = z - \frac{\varepsilon_3}{z}, \quad (118)$$

$$z = \sqrt[3]{\left(-\frac{\varepsilon_1}{2}\right) \pm \sqrt{\varepsilon_2}}, \quad (119)$$

$$\varepsilon_1 = \frac{1}{y^2}. \quad (120)$$

$$\text{The discriminant of the cubic equation } \varepsilon_2 = \frac{1}{4y^2} + \frac{(\varepsilon_3)^3}{27}, \quad (121)$$

$$\varepsilon_3 \text{ is given by } \varepsilon_3 = -3 - \frac{1}{y^2} \quad (122)$$

After some lengthy calculations, it can be verified that:

$$\rho = 1 \mp \left((z - 1) + \frac{\left(3 + \frac{1}{y^2}\right)^{\frac{1}{2}}}{z} \right)^{\frac{1}{2}}, \quad (123)$$

$$\text{where } z = \sqrt[3]{-\frac{1}{2y^2} \pm i \sqrt{\frac{1}{12y^4} + 1 + \frac{1}{2y^2} + \frac{1}{27y^6}}}, i = \sqrt{-1}.$$

By Eq. (102), we have:

$$1 \mp \left((z - 1) + \frac{\left(3 + \frac{1}{\rho^2}\right)^{\frac{1}{2}}}{z} \right)^{\frac{1}{2}} = \varphi_1^{-1}(\rho), \quad (124)$$

$$\text{where } z = \sqrt[3]{-\frac{1}{2\rho^2} \pm i \sqrt{\frac{1}{12\rho^4} + 1 + \frac{1}{2\rho^2} + \frac{1}{27\rho^6}}}, i = \sqrt{-1}.$$

This completes the proof.

VI. To prove VI), it is enough to show that for all $\beta_1, \beta_2 \in (1, \infty)$ such that $\varphi_2(\beta_1) = \varphi_2(\beta_2)$, then β_1 and β_2 should never be distinct.

Let $\varphi_2(\beta_1) = \varphi_2(\beta_2)$. Then $\frac{(1+\beta_1)}{(1+((1+\beta_1)^2)^{\frac{2}{3}})} = \frac{(1+\beta_2)}{(1+((1+\beta_2)^2)^{\frac{2}{3}})}$, such that $\beta_1, \beta_2 \in (1, \infty), \beta_1 \neq \beta_2$. This implies

Hence,

$$(\beta_2 - \beta_1)(\beta_1 + \beta_2 + 2)[(1 + \beta_1)^2(1 + \beta_2)^2((1 + \beta_1)^2 + (1 + \beta_2)^2 + 2)] = 0. \quad (125)$$

Eq. (125) generates three possible cases:

$$\text{Case 1. } \beta_2 - \beta_1 = 0. \text{ Hence } \rho_1 = \rho_2 \text{ (contradiction to } \beta_1 \neq \beta_2). \quad (126)$$

$$\text{Case 2. } \beta_1 + \beta_2 + 2 = 0. \text{ Hence } \beta_1 + \beta_2 = -2 \text{ (contradiction, since M/G/1 is a stable QM, } \beta_1, \beta_2 \in (1, \infty)). \quad (127)$$

$$\text{Case 3. } [(1 + \beta_1)^2(1 + \beta_2)^2((1 + \beta_1)^2 + (1 + \beta_2)^2 + 2)] = 0. \quad (128)$$

By $\beta_1, \beta_2 \in (1, \infty)$, following mathematical analysis,

$$[(1 + \beta_1)^2(1 + \beta_2)^2((1 + \beta_1)^2 + (1 + \beta_2)^2 + 2)] > 96, \text{ which directly implies by Eq. (126).}$$

$$0 > 96 \text{ (contradiction)}. \quad (129)$$

Based on the above analysis, VI) follows.

VII. It suffices to show that for all $\beta_1, \beta_2 \in (1, \infty)$ such that $\varphi_2(\beta_1) = \varphi_2(\beta_2)$, then $\beta_1 = \beta_2$ holds. The proof is clearly immediate from Eq. (126).

VIII. Since M/G/1 QM is stable, the condition $\beta > 1$ and a similar proof to that in II), VIII) follows.

The proof of x) is analogous to III).

IX. We have $\varphi_1^{-1}(\rho) = 1 \mp \left((z-1) + \frac{(3+\frac{1}{\rho^2})}{z} \right)^{\frac{1}{2}}$, for $\rho \in (0,1)$, Eq. (103).

Hence,

$$|1 - \varphi_1^{-1}(\rho)|^2 = \left| (z-1) + \frac{(3+\frac{1}{\rho^2})}{z} \right| \leq |z| + 1 + \frac{(3+\frac{1}{\rho^2})}{|z|} \quad (130)$$

By Eq. (124),

$$z = \sqrt[3]{-\frac{1}{2\rho^2} \pm i \sqrt{\frac{1}{12\rho^4} + 1 + \frac{1}{2\rho^2} + \frac{1}{27\rho^6}}}$$

This implies:

$$|z|^6 = \left(\frac{1}{4\rho^4} + \frac{1}{12\rho^4} + 1 + \frac{1}{2\rho^2} + \frac{1}{27\rho^6} \right) < \left(\frac{1}{4\rho^6} + \frac{1}{12\rho^6} + \frac{1}{\rho^6} + \frac{1}{2\rho^6} + \frac{1}{27\rho^6} \right) = \frac{101}{54\rho^6}, \text{ or } |z| < \frac{(\frac{101}{54})^{\frac{1}{6}}}{\rho} < \frac{(\frac{101}{54})^{\frac{1}{6}}}{\rho^2}. \quad (131)$$

Moreover, by the above step, it is clear that:

$$|z|^6 = \left(\frac{1}{4\rho^4} + \frac{1}{12\rho^4} + 1 + \frac{1}{2\rho^2} + \frac{1}{27\rho^6} \right) > 1, \text{ which directly implies } |z| > 1 \left(\text{equivaently, } \frac{1}{|z|} < 1 \right). \quad (132)$$

Thus, it is obtained that

$$|1 - \varphi_1^{-1}(\rho)|^2 < \frac{(\frac{101}{54})^{\frac{1}{6}}}{\rho^2} + 2 + \left(3 + \frac{1}{\rho^2} \right) = 5 + \frac{(1 + (\frac{101}{54})^{\frac{1}{6}})}{\rho^2}.$$

This completes the proof.

X. The stability of the underlying QM implies $\beta > 1$. We have $\varphi_2^{-1}(\beta) = -1 \mp \left((z-1) + \frac{(3+\frac{1}{\beta^2})}{z} \right)^{\frac{1}{2}}$

Hence,

$$|1 + \varphi_2^{-1}(\beta)|^2 = \left| (z-1) + \frac{(3+\frac{1}{\beta^2})}{z} \right| \leq |z| + 1 + \frac{(3+\frac{1}{\beta^2})}{|z|}. \quad (133)$$

Hence,

$$z = \sqrt[3]{-\frac{1}{2\beta^2} \pm i \sqrt{\frac{1}{12\beta^4} + 1 + \frac{1}{2\beta^2} + \frac{1}{27\beta^6}}, i = \sqrt{-1}}.$$

So, $|z| < \left(\frac{101}{54}\right)^{\frac{1}{6}} Eq.$ (132). Also, it can be verified that $\frac{1}{|z|} < \frac{\beta^2}{\left(\frac{101}{54}\right)^{\frac{1}{6}}} Eq.$ (133). Consequently, (XIII) will follow.

xiv) we have $\frac{\partial \varphi_1}{\partial \rho} = \frac{2(1-\rho)^2-1}{(1+(1-\rho)^2)^{\frac{5}{2}}}$. Hence, $\frac{\partial \varphi_1}{\partial \rho} > 0 (< 0)$ if and only if $(1-\rho)^2 > \frac{1}{2} (< \frac{1}{2})$. By (PT) 2.15, φ_1 is increasing (decreasing) in ρ if and only if $(1-\rho)^2 > \frac{1}{2} (< \frac{1}{2})$. According to the stability of M/G/1 QM, $\rho \in (0,1)$. Hence, it follows that $(1-\rho)^2 \in (0,1)$. Consequently, xiv) follows:

$$\varphi_2(\beta) = K_2 = \frac{-(1+\beta)}{(1+(1+\beta)^2)^{\frac{3}{2}}}$$

xv) We have $\frac{\partial \varphi_2}{\partial \beta} = \frac{2(1+\beta)^2-1}{(1+(1+\beta)^2)^{\frac{5}{2}}}$. Hence, $\frac{\partial \varphi_2}{\partial \beta} > 0 (< 0)$ if and only if $(1+\beta)^2 > \frac{1}{2} (< \frac{1}{2})$. By (PT), φ_2 is increasing (decreasing) in ρ if and only if $(1+\beta)^2 > \frac{1}{2} (< \frac{1}{2})$. According to the stability of M/G/1 QM, $\beta \in (1, \infty)$. Consequently, xv) follows.

7 | $e^{\text{FIM}(M/G/1)}$ and Impact of Stability of M/G/1 Qm on the Stability of Fim

7.1 | Exponential Matrix of FIM

Theorem 10. $e^{\text{FIM}(M/G/1)}$ solves $\frac{dx}{dt} = Ax$.

Proof: It is shown that $[g_{ij}]$ of Theorem 3 is:

$$[g_{ij}] = \begin{pmatrix} \frac{1}{(1-\rho)^2} & 0 \\ 0 & \frac{-1}{(\beta+1)^2} \end{pmatrix}. \quad (134)$$

We write

$$[g_{ij}] = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, a = \left(\frac{1}{(1-\rho)^2}\right), b = \frac{-1}{(\beta+1)^2}. \quad (135)$$

Thus,

$$\Phi(\delta) = \det \begin{pmatrix} a-\delta & 0 \\ 0 & b-\delta \end{pmatrix} = 0.$$

Therefore,

$$\delta^2 - (a+b)\delta + ab = 0, \text{ so } \delta_{1,2} = a, b.$$

Hence,

$$D = \begin{pmatrix} \delta_1 & 0 \\ 0 & \delta_2 \end{pmatrix}, \quad (136)$$

For $\delta_{1,2} = a, b$

$$\text{Hence, } T = T^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (137)$$

Thus, $e^{\text{FIM}(M/G/1)}$ reads as:

$$e^{\text{FIM}(M/G/1)} = Te^DT^{-1} = \begin{pmatrix} e^a & 0 \\ 0 & e^b \end{pmatrix}. \quad (138)$$

This proves that IME of the underlying QM solves the following:

$$\frac{dx}{dt} = Ax. \quad (139)$$

7.2 | Impact of Stability of M/G/1 QM on the Stability of FIM

Theorem 11. The stability of FIM of the underlying QM holds \Leftrightarrow the underlying QM is unstable.

Proof: Following the Preliminary Theorem, it suffices to show that:

FIM's eigenvalues of FIM of the underlying QM are negative real numbers \Leftrightarrow the instability of the underlying QM is satisfied.

It holds by of *Theorem 10* that the eigenvalues of FIM are $\delta_{1,2} = a, b$, $a = \left(\frac{1}{(1-\rho)^2}\right)$, $b = \frac{-1}{(\beta+1)^2}$. clearly, $b = \frac{-1}{(\beta+1)^2} < 0$. Therefore, the proof would be immediate if we proved that:

$$a = \left(\frac{1}{(1-\rho)^2}\right) < 0 \Leftrightarrow \text{the instability of the underlying QM is satisfied.}$$

We first prove the necessity condition, $a = \left(\frac{1}{(1-\rho)^2}\right) < 0 \Rightarrow$ M/G/1 QM is unstable. Assume that FIM is stable, then $a = \left(\frac{1}{(1-\rho)^2}\right) < 0$ follows. This implies $\frac{1}{(1-\rho)} = im$, $i = \sqrt{-1}$, m is any real number. Hence, $(1-\rho) = -im$. Consequently, $\rho = 1 + im$, $|\rho| = \sqrt{1+m^2} > 1$. In other words, M/G/1 QM is unstable.

To prove sufficiency, let M/G/1 QM be unstable. Then, $\rho > 1$. This directly implies, $|\rho| > 1$. This rewrites ρ to be of the form $\rho = 1 + im$, m is any real number. Clearly, this implies, $(1-\rho) = -im$, or $\frac{1}{(1-\rho)} = im$. Thus, it holds that $a = \left(\frac{1}{(1-\rho)^2}\right) = -\frac{1}{m^2} < 0$, which proves that FIM is stable.

7.3 | Revealing Queue-Fisher Information Matrix Unifiers (QFIMU)

Throughout this section, we introduce QFIMU, to be devised by the function:

$$\eta(\rho, \beta) = [g_{ij}] \begin{pmatrix} \rho \\ -\beta \end{pmatrix} = \begin{pmatrix} \frac{1}{(1-\rho)^2} & 0 \\ 0 & \frac{-1}{(\beta+1)^2} \end{pmatrix} \begin{pmatrix} \rho \\ -\beta \end{pmatrix} = \begin{pmatrix} \frac{\rho}{(1-\rho)^2} \\ \frac{-\beta}{(\beta+1)^2} \end{pmatrix}. \quad (140)$$

where $\rho = 1 - p(0)$ and $\beta = C_s^2$.

Theorem 12. The function η Eq. (140) satisfies the following:

- I. η is well-defined.
- II. η is One-to-One.
- III. η is surjective.
- IV. η has a unique inverse, η^{-1} determined, which is characterized by

$$\left| \eta^{-1}(\rho) - \left(1 + \frac{1}{2\rho}\right) \right|^2 < \frac{2}{\rho^2}. \quad (141)$$

and

$$\left| \eta^{-1}(\beta) + \left(1 + \frac{1}{2\beta}\right) \right|^2 < 2. \quad (142)$$

Proof: to prove I), it suffices to show that for all (ρ_1, β_1) , (ρ_2, β_2) such that $\rho_1 \neq \rho_2$ and $\beta_1 \neq \beta_2$ and $\eta(\rho_1, \beta_1) = \eta(\rho_2, \beta_2)$.

$$(143)$$

By Eq. (143), we have

$$\frac{\rho_1}{(1 - \rho_1)^2} = \frac{\rho_2}{(1 - \rho_2)^2} \quad (144)$$

and

$$\frac{\beta_1}{(1 + \beta_1)^2} = \frac{\beta_2}{(1 + \beta_2)^2} \quad (145)$$

By Eq. (144), one gets:

$$(\rho_1 - \rho_2)(1 + \rho_1\rho_2) = 0. \quad (146)$$

Eq. (146) implies either $\rho_1 = \rho_2$ or $\rho_2 = \frac{-1}{\rho_1}$ (contradiction, since, for example, if $\rho_1 = 2$ implies that $\rho_2 = -0.5 \notin (0,1)$, i. e., enforcing instability of the underlying stable M/G/1 QM). Therefore, the ρ branch of the QIFMU is well-defined.

Following Eq. (145), we have:

$$(\beta_1 - \beta_2)(1 - \beta_1\beta_2) = 0. \quad (147)$$

Eq. (147) implies either $\beta_1 = \beta_2$ or $\beta_2 = \frac{1}{\beta_1}$ (contradiction, since, for example, if $\beta_1 = 2$ implies that $\beta_2 = 0.5 \notin (1, \infty)$, i. e., enforcing instability of the underlying stable M/G/1 QM). Therefore, the β branch of the QIFMU is well-defined. This completes the proof of I).

As for II), it suffices to show that all $(\rho_1, \beta_1), (\rho_2, \beta_2)$ such that:

$$\eta(\rho_1, \beta_1) = \eta(\rho_2, \beta_2) \text{ implies } (\rho_1, \beta_1) = (\rho_2, \beta_2). \quad (148)$$

Clearly, by Eqs. (136)-(138) is satisfied. Hence, II) follows.

Clearly, by Eq. (140), both of $\frac{\rho}{(1-\rho)^2}$ and $\frac{-\beta}{(\beta+1)^2}$ are uniquely characterized by ρ and β , respectively. Thus, III) holds.

iv) To compute η^{-1} , assume that there exists x, y such that

$$\eta(\rho, \beta) = \begin{pmatrix} \frac{1}{(1-\rho)^2} & 0 \\ 0 & \frac{-1}{(\beta+1)^2} \end{pmatrix} \begin{pmatrix} \rho \\ -\beta \end{pmatrix} = \begin{pmatrix} \frac{\rho}{(1-\rho)^2} \\ \frac{-\beta}{(\beta+1)^2} \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}. \quad (149)$$

Therefore,

$$\frac{\rho}{(1 - \rho)^2} = x. \quad (150)$$

and

$$\frac{-\beta}{(\beta + 1)^2} = y. \quad (151)$$

Following Eq. (148), one gets:

$$\rho = \frac{(2 + \frac{1}{x}) \pm \frac{1}{x} \sqrt{1 + 4x^2}}{2}. \quad (152)$$

Using Eq. (149) and following a similar argument as in Eq. (150), we have:

$$\beta = \frac{-(2 + \frac{1}{y}) \pm \frac{1}{y} \sqrt{1 + 4y^2}}{2}. \quad (153)$$

Based on Eq. (141) and Eq. (142), it is determined for both ρ and branches of η^{-1} would respectively satisfy that

$$|\eta^{-1}(\rho) - (1 + \frac{1}{2\rho})|^2 = \frac{(1+4\rho^2)}{4\rho^2} = 1 + \frac{1}{4\rho^2} < \frac{1}{\rho^2} + \frac{1}{\rho^2} = \frac{2}{\rho^2} \text{ (since } \rho \in (0,1)\text{)}. \quad (141)$$

Following a similar argument, it could be shown that

$$|\eta^{-1}(\beta) + (1 + \frac{1}{2\beta})|^2 = \frac{(1+4\beta^2)}{4\beta^2} = 1 + \frac{1}{4\beta^2} < 1 + 1 = 2 \text{ (since } \beta \in (1,\infty)\text{)}. \quad (142)$$

This completes the proof of our theorem.

8 | RICCI Scalar, \mathcal{R} , Curvature of Space Time (Einestein Tensor) \wp

Stress Energy Tensor R, Ω , the corresponding threshold theorems for the underlying curvatures and the dual queueing impact on the existence of the Inverse Fisher Information Matrix (IFIM).

Theorem 13. The underlying QM satisfies:

- I. The Ricci scalar subject to Angular Technique, \mathcal{R}_{AT} is determined by.

$$\mathcal{R}_{AT} = \frac{2(\rho - 1)(1 + \beta)}{(1 + (1 - \rho)^2)^{\frac{3}{2}}(1 + (1 + \beta)^2)^{\frac{3}{2}}}, \quad (143)$$

where $\rho = 1 - p(0)$ and C_s^2 define server utilization and Squared coefficient of variations, respectively.

- II. $\mathcal{R}_{AT} \rightarrow 0$ if and only if $\rho = 1$.

- III. $\mathcal{R}_{AT} \rightarrow 0$ if and only if $\beta \rightarrow \infty$.

for all $\rho \neq 1$ (equivalently, whether the underlying QM is either stable or unstable)

- IV. M/G/1 QM is unstable \Leftrightarrow

There exists a small enough positive number ϵ , with $\epsilon \rightarrow 0$ such that $A_{\text{curved}}(\epsilon), A_{\text{flat}}(\epsilon)$ Eq. (43) must satisfy:

$$A_{\text{curved}}(\epsilon) \gtrsim A_{\text{flat}}(\epsilon). \quad (144)$$

- V. The Spacetime curvature (Einstein Tensor) subject to Angular Technique, \wp_{AT} is determined by

$$\wp_{AT} = \begin{pmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{pmatrix}, \quad (145)$$

where the components G_{11}, G_{12}, G_{21} and G_{22} are determined by

$$G_{11} = \frac{(1 + \beta)}{(1 - \rho)(1 + (1 - \rho)^2)^{\frac{3}{2}}(1 + (1 + \beta)^2)^{\frac{3}{2}}}, \quad (146)$$

$$G_{12} = 0, \quad (147)$$

$$G_{21} = \frac{(\alpha - 1)}{(1 - \rho)^2} \quad (148)$$

where α is the curvature parameter.

$$G_{22} = \frac{(\rho - 1)}{(1 + \beta)(1 + (1 - \rho)^2)^{\frac{3}{2}}(1 + (1 + \beta)^2)^{\frac{3}{2}}}. \quad (149)$$

- VI. The stress-energy tensor ϖ is devised by

$$\Omega = \begin{pmatrix} \varpi_{11} & \varpi_{12} \\ \varpi_{21} & \varpi_{22} \end{pmatrix}, \quad (150)$$

where the components ϖ_{11} , ϖ_{12} , ϖ_{21} and ϖ_{22} are determined by

$$\varpi_{11} = \frac{c^4 G_{11}}{8\pi g}, \quad (151)$$

$$\varpi_{12} = 0,$$

$$\varpi_{21} = \frac{c^4 G_{21}}{8\pi g}, \quad (152)$$

$$\varpi_{22} = \frac{c^4 G_{22}}{8\pi g}, \quad (153)$$

where g is the universal gravitational constant, c is the speed of light:

VII.

$$\mathcal{R}_{AT} = \begin{cases} \text{increasing in } \rho, & \rho = 1 + \frac{m}{\sqrt{2}} \text{ (instability phase),} \\ \text{increasing in } \rho, & \rho = 1 - \frac{m}{\sqrt{2}} \text{ (stability phase).} \end{cases} \quad (154)$$

Provided that $m > 1$.

VIII.

$$\mathcal{R}_{AT} = \begin{cases} \text{decreasing in } \rho, & \rho = 1 + \frac{m}{\sqrt{2}} \text{ (instability phase),} \\ \text{decreasing in } \rho, & \rho = 1 - \frac{m}{\sqrt{2}} \text{ (instability phase).} \end{cases} \quad (155)$$

Provided that $1 > m > 0$.

$$\text{X. } \mathcal{R}_{AT} = \begin{cases} \text{decreasing in } \beta, & \text{the underlying QM is stable,} \\ \text{increasing in } \beta, & \text{the underlying QM is unstable.} \end{cases} \quad (156)$$

$$\text{XI. } G_{11} = \begin{cases} \text{increasing in } \rho, & \text{the underlying QM is stable or unstable, } \rho \neq 1, \\ \text{decreasing in } \rho, & \text{the underlying QM is stable.} \end{cases} \quad (157)$$

XII. G_{21} is forever decreasing in α (curvature parameter) whether M/G/1 QM is stable or unstable. If $\rho = 1$, the decreasability of G_{21} in α is undecidable.

XIII. G_{21} is forever increasing (decreasing) in ρ if M/G/1 QM is stable, $\alpha < 1$ ($\alpha > 1$).

XV. G_{21} is forever decreasing in ρ if either one of the following branches holds:

$$\begin{cases} \rho \in (0,1), & \text{the underlying QM is stable,} & \alpha > 1, \\ \rho > 1, & \text{the underlying QM is unstable,} & \alpha < 1. \end{cases} \quad (158)$$

XVI. G_{22} is forever increasing in ρ .

XVII. G_{22} is forever increasing (decreasing) in β if and only if $\rho < 1$ ($\rho > 1$). If $\rho = 1$, the decision is undecidable.

Proof:

I. Immediate.

II. By I), $\mathcal{R}_{AT} = \frac{2(\rho-1)(1+\beta)}{(1+(1-\rho)^2)^{\frac{3}{2}}(1+(1+\beta)^2)^{\frac{3}{2}}}$. Hence, $\mathcal{R}_{AT} \rightarrow 0$ if and only if $(\rho - 1)(1 + \beta) \rightarrow 0$. Since $\beta > 1$, the required result follows.

III. Since M/G/1 is a stable QM, $\rho \in (0,1)$ holds. This implies for all $\rho \neq 1$ (equivalently, whether the underlying QM is either stable or unstable) $\mathcal{R}_{AT} \rightarrow 0$ if and only if $\frac{2(\rho-1)}{(1+(1-\rho)^2)^{\frac{3}{2}}} \lim_{\beta \rightarrow \infty} \frac{(1+\beta)}{(1+(1+\beta)^2)^{\frac{3}{2}}} = 0$. By $(1 + (1 + \beta)^2)^{\frac{3}{2}} > (1 + \beta)$, the proof follows.

I. M/G/1 QM is unstable if and only if $\rho \geq 1$, or equivalently $\mathcal{R}_{AT} \geq 0$. This holds if and only if

$$\mathcal{R}_{AT} = \lim_{\epsilon \rightarrow 0} \frac{6n}{\epsilon^2} \left[1 - \frac{A_{\text{curved}}(\epsilon)}{A_{\text{flat}}(\epsilon)} \right] \geq 0. \quad (159)$$

This is equivalent to $\left[1 - \frac{A_{\text{curved}}(\epsilon)}{A_{\text{flat}}(\epsilon)} \right] \geq 0$. This completes the proof.

By Eq. (44), $G_{ij} = R_{ij}^{(\alpha)} \frac{\mathcal{R}_{AT}}{2} \mathbf{g}_{ij} = \frac{8\pi\varrho\omega_{ij}}{c^4}$, $i, j = 1, 2$. Hence, it follows that:

$$G_{11} = R_{11}^{(\alpha)} \frac{\mathcal{R}_{AT}}{2} \mathbf{g}_{11} = \frac{8\pi\varrho\omega_{11}}{c^4}, \quad (160)$$

$$G_{12} = R_{12}^{(\alpha)} \frac{\mathcal{R}_{AT}}{2} \mathbf{g}_{12} = \frac{8\pi\varrho\omega_{12}}{c^4}, \quad (161)$$

$$G_{21} = R_{21}^{(\alpha)} \frac{\mathcal{R}_{AT}}{2} \mathbf{g}_{11} = \frac{8\pi\varrho\omega_{21}}{c^4}, \quad (162)$$

$$G_{22} = R_{22}^{(\alpha)} \frac{\mathcal{R}_{AT}}{2} \mathbf{g}_{22} = \frac{8\pi\varrho\omega_{22}}{c^4}. \quad (163)$$

Using Theorems 3 and 8, together with Eqs. (160)-(163), the proof of V and VI will follow.

VII. It could be verified that:

$$\frac{\partial \mathcal{R}_{AT}}{\partial \rho} = \frac{2(1 - 2(1 - \rho)^2)(1 + \beta)}{(1 + (1 - \rho)^2)^{\frac{5}{2}}(1 + (1 + \beta)^2)^{\frac{3}{2}}}. \quad (164)$$

Therefore,

$$\frac{\partial \mathcal{R}_{AT}}{\partial \rho} < 0 (> 0) \text{ if and only if } (1 - 2(1 - \rho)^2) < 0 (> 0). \quad (165)$$

if $(1 - 2(1 - \rho)^2) > 0$ is satisfied $\Leftrightarrow \exists m \in (1, \infty)$ satisfying $(1 - \rho)^2 = \frac{m^2}{2}$, which $1 - \rho = \pm \frac{m}{\sqrt{2}}$ or $\rho = 1 \mp \frac{m}{\sqrt{2}}$. Following the Preliminary Theorem (PT), this implies \mathcal{R}_{AT} is increasing in ρ if and only if $\rho = 1 \mp \frac{m}{\sqrt{2}}$. For or $\rho = 1 + \frac{m}{\sqrt{2}}$, this enforces or $\rho > 1$, which violates the underlying QM's stability, or $\rho = 1 - \frac{m}{\sqrt{2}}$, this enforces $\rho < 1$, which guarantees the stability of M/G/1 QM.

On the other hand, if $(1 - 2(1 - \rho)^2) < 0$, it holds by PT, that \mathcal{R}_{AT} is decreasing in ρ if and only if there exists $m \in (0,1)$ satisfying $(1 - \rho)^2 = \frac{m^2}{2}$, which $1 - \rho = \pm \frac{m}{\sqrt{2}}$. For or $\rho = 1 + \frac{m}{\sqrt{2}}$, this enforces or $\rho > 1$, which indicates the underlying QM's instability. Moreover, or $\rho = 1 - \frac{m}{\sqrt{2}}$, this enforces $\rho > 1 - \frac{1}{\sqrt{2}}$, which violates the stability of M/G/1 QM.

Following the above analytic results, VII) and VIII) are immediate.

X. It could be shown that

$$\frac{\partial \mathcal{R}_{AT}}{\partial \beta} = \frac{2(2(1 + \beta)^2 - 1)(1 - \rho)}{(1 + (1 - \rho)^2)^{\frac{3}{2}}(1 + (1 + \beta)^2)^{\frac{5}{2}}}. \quad (166)$$

Undertaking a similar mathematical mechanism as in VII) and VIII), the proof follows.

XI. After some mathematical manipulation, we have

$$\frac{\partial G_{11}}{\partial \rho} = \frac{(2(1 - \rho)^2 + 1)(1 - \rho)(1 + \beta)}{(1 - \rho)^2(1 + (1 - \rho)^2)^{\frac{5}{2}}(1 + (1 + \beta)^2)^{\frac{3}{2}}}. \quad (167)$$

Eq. (167) provides an evidence that $\frac{\partial G_{11}}{\partial \rho} > 0$ for all $\rho \neq 1$. Applying the Preliminary Theorem shows that G_{11} is forever increasing in ρ . This is applicable for stable and unstable M/G/1 QM, with $\rho \neq 1$ (since $\rho = 1$, violates the continuity requirement of $\frac{\partial G_{11}}{\partial \rho}$). Furthermore, it could be proved that G_{11} is increasing in ρ when the underlying QM is in the stability phase.

Moreover,

$$\frac{\partial G_{11}}{\partial \beta} = \frac{(1 - 2(1 + \beta)^2)(1 - \rho)(1 + \beta)}{(1 - \rho)(1 + (1 - \rho)^2)^{\frac{3}{2}}(1 + (1 + \beta)^2)^{\frac{5}{2}}}. \quad (168)$$

Clearly, from Eq. (168) and PT Eq. (14), it follows that G_{11} is never increasing in β (since $\rho = 1$, violates the continuity requirement of $\frac{\partial G_{11}}{\partial \beta}$). Let us assume that G_{11} is never increasing in β . This implies:

$$(1 - 2(1 + \beta)^2) > 0. \quad (169)$$

This is equivalent to $(1 + \beta)^2 < \frac{1}{2}$ (contradiction, since the stability of M/G/1 QM enforces the requirements $\rho \in (0,1)$ and $\beta \in (1, \infty)$). This suggests that the only left possibility is that G_{11} is forever increasing in β , or equivalently by Preliminary Theorem, to $(1 + \beta)^2 > \frac{1}{2}$, a satisfied condition by stable M/G/1 QM.

XII. $\frac{\partial G_{21}}{\partial \alpha} = \frac{-1}{(1-\rho)^2}$, implies by the preliminary theorem, that G_{21} is forever decreasing in α whether M/G/1 QM is stable or unstable. If $\rho = 1$, the decreasability of G_{21} in α is undecidable, since this means $\frac{\partial G_{21}}{\partial \alpha} \rightarrow \infty$.

Furthermore, $\frac{\partial G_{21}}{\partial \rho} = \frac{2(1-\alpha)}{(1-\rho)^3}$. Consequently, $\frac{\partial G_{21}}{\partial \rho} > 0$ if and only if M/G/1 QM is stable, $\alpha < 1$, $\frac{\partial G_{21}}{\partial \rho} < 0$ if and only if M/G/1 QM is stable, $\alpha > 1$. By the Preliminary Theorem, it follows that G_{21} is forever increasing(decreasing) in ρ if M/G/1 QM is stable, $\alpha < 1$ ($\alpha > 1$). This proves XIII).

Engaging the same technique proves xv).

We have:

$$\frac{\partial G_{22}}{\partial \rho} = \frac{(4(1-\rho)^2+1)}{(1+\beta)(1+(1-\rho)^2)^{\frac{5}{2}}(1+(1+\beta)^2)^{\frac{3}{2}}}, \text{ and } \frac{\partial G_{22}}{\partial \beta} = \frac{(3(1+\beta)^2+1)(1-\rho)}{(1+\beta)^2(1+(1-\rho)^2)^{\frac{5}{2}}(1+(1+\beta)^2)^{\frac{3}{2}}}. \quad (170)$$

Engaging our technique, the reader can easily verify that both XVI) and XVII) will hold. This completes the proof of our Theorem.

Theorem 14. The underlying QM satisfies:

I. Ricci scalar subject to Monge Technique, \mathcal{R}_{MT} is determined by:

$$\mathcal{R}_{MT} = -2. \quad (171)$$

II. There exists a small enough positive number ϵ , with $\epsilon \rightarrow 0$ such that $A_{\text{curved}}(\epsilon)$, $A_{\text{flat}}(\epsilon)$ Eq. (43) must satisfy.

$$A_{\text{curved}}(\epsilon) \rightarrow (A_{\text{flat}}(\epsilon)), \quad (172)$$

III. The Spacetime curvature (Einstein Tensor) subject to Angular Technique, \wp_{AT} is determined by

$$\wp_{MT} = \begin{pmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{pmatrix}, \quad (173)$$

where the components G_{11} , G_{12} , G_{21} and G_{22} are determined by

$$G_{11} = \frac{(1 - \alpha)}{(1 - \rho)^2} = G_{21}, \quad (174)$$

where α is the curvature parameter.

$$G_{12} = 0, \quad (175)$$

$$G_{22} = -\frac{1}{(1+\beta)^2}. \quad (176)$$

IV. The stress-energy tensor ϖ is devised by

$$\varpi_{MT} = \begin{pmatrix} \varpi_{11} & \varpi_{12} \\ \varpi_{21} & \varpi_{22} \end{pmatrix}, \quad (177)$$

where the components $\varpi_{11}, \varpi_{12}, \varpi_{21}$ and ϖ_{22} are determined by

$$\varpi_{11} = \frac{c^4 G_{11}}{8\pi\mathcal{G}}, \quad (178)$$

$$\varpi_{12} = 0, \quad (179)$$

$$\varpi_{21} = \frac{c^4 G_{21}}{8\pi\mathcal{G}}, \quad (180)$$

$$\varpi_{22} = \frac{c^4 G_{22}}{8\pi\mathcal{G}}, \quad (181)$$

where \mathcal{G} is the universal gravitational constant, c is the speed of light.

V.

$$G_{11} = G_{21} = \begin{cases} \text{increasing in } \rho, & \text{the underlying QM is stable or unstable,} \\ \text{decreasing in } \beta, & \text{the underlying QM is stable.} \end{cases} \quad (182)$$

VI. G_{11} is forever decreasing in α (curvature parameter) whether M/G/1 QM is stable or unstable. If $\rho = 1$, the decreasability of G_{21} in α is undecidable.

VII. G_{11} is forever increasing(decreasing) in ρ if M/G/1 QM is stable, $\alpha < 1$ ($\alpha > 1$).

VIII. G_{11} is forever decreasing in ρ if either one of the following branches holds:

$$\begin{cases} \rho \in (0,1), & \text{the underlying QM is stable,} & \alpha > 1, \\ \rho > 1, & \text{the underlying QM is unstable,} & \alpha < 1. \end{cases} \quad (183)$$

X. G_{22} is forever increasing in β .

XI. G_{22} is forever increasing (decreasing) in β if and only if $\rho < 1$ ($\rho > 1$). If $\rho = 1$, the decision is undecidable.

Proof:

I. We have $K_{G,MT} = -1$. We have $\mathcal{R}_{MT} = 2K_{G,MT} = -2$, Eq. (171).

II. Since $\mathcal{R}_{MT} = -2$. Hence,

$$\mathcal{R}_{MT} = \lim_{\epsilon \rightarrow 0} \frac{6n}{\epsilon^2} \left[1 - \frac{A_{\text{curved}}(\epsilon)}{A_{\text{flat}}(\epsilon)} \right] = -2. \quad (184)$$

This is equivalent to $\left[1 - \frac{A_{\text{curved}}(\epsilon)}{A_{\text{flat}}(\epsilon)} \right] \gtrsim -2$. This completes the proof.

By Eq. (189), $G_{ij} = R_{ij}^{(\alpha)} \frac{\mathcal{R}_{MT}}{2} \mathbf{g}_{ij} = \frac{8\pi\mathcal{G}\varpi_{ij}}{c^4}$, $i, j = 1, 2$. Hence, it follows that:

$$G_{11} = R_{11}^{(\alpha)} \frac{\mathcal{R}_{MT}}{2} \mathbf{g}_{11} = \frac{8\pi\mathcal{G}\varpi_{11}}{c^4}, \quad (185)$$

$$G_{12} = R_{12}^{(\alpha)} \frac{\mathcal{R}_{MT}}{2} \mathbf{g}_{12} = \frac{8\pi\mathcal{G}\varpi_{12}}{c^4}, \quad (186)$$

$$G_{21} = R_{21}^{(\alpha)} \frac{\mathcal{R}_{MT}}{2} \mathbf{g}_{11} = \frac{8\pi\mathcal{G}\varpi_{21}}{c^4}, \quad (187)$$

$$G_{22} = R_{22}^{(\alpha)} \frac{\mathcal{R}_{MT}}{2} \mathbf{g}_{22} = \frac{8\pi\mathcal{G}\varpi_{22}}{c^4}. \quad (188)$$

Using *Theorems 3* and *8*, *Eq. (171)* together with *Eqs. (185)-(188)*, the proof of III and IV will follow.

The remaining proofs of V-XI are omitted since they are provable by following the same analytic mechanism undertaken in *Eqs. (154)-(158)*.

Theorem 15. The underlying QM satisfies:

- I. \mathcal{R}_{AT} has a relative minimum of $(1 \mp \frac{1}{\sqrt{2}}, -1 \pm \frac{1}{\sqrt{2}})$.
- II. Both maxima and minima for all the components of the Spacetime curvature (Einstein Tensor) subject to Angular Technique, \wp_{AT} is undecidable.

Proof:

I. We have $\frac{\partial \mathcal{R}_{AT}}{\partial \rho} = \frac{2(1-2(1-\rho)^2)(1+\beta)}{(1+(1-\rho)^2)^{\frac{5}{2}}(1+(1+\beta)^2)^{\frac{3}{2}}}$, $\frac{\partial \mathcal{R}_{AT}}{\partial \beta} = \frac{2(2(1+\beta)^2-1)(1-\rho)}{(1+(1-\rho)^2)^{\frac{3}{2}}(1+(1+\beta)^2)^{\frac{5}{2}}}$ *Eqs. (164)* and *(166)* respectively.

Hence, $\frac{\partial \mathcal{R}_{AT}}{\partial \rho} = 0 = \frac{\partial \mathcal{R}_{AT}}{\partial \beta}$. The critical points are $(\rho_{critical}, \beta_{critical}) = (1 \mp \frac{1}{\sqrt{2}}, -1 \pm \frac{1}{\sqrt{2}})$. Moreover, we have:

$$\frac{\partial^2 \mathcal{R}_{AT}}{\partial \rho^2} = \frac{6(3-2(1-\rho)^2)(1+\beta)}{(1+(1-\rho)^2)^{\frac{7}{2}}(1+(1+\beta)^2)^{\frac{3}{2}}}, \frac{\partial^2 \mathcal{R}_{AT}}{\partial \beta^2} = \frac{2(1-14(1+\beta)^2)(\rho-1)}{(1+(1-\rho)^2)^{\frac{3}{2}}(1+(1+\beta)^2)^{\frac{7}{2}}}, \frac{\partial^2 \mathcal{R}_{AT}}{\partial \rho \beta} = \frac{2(1-2(1-\rho)^2)(1-2(1+\beta)^2)}{(1+(1-\rho)^2)^{\frac{5}{2}}(1+(1+\beta)^2)^{\frac{5}{2}}}. \tag{189}$$

Hence, by *Eq. (45)*:

$$D\left(1 - \frac{1}{\sqrt{2}}, -1 + \frac{1}{\sqrt{2}}\right) = \frac{\partial^2 \mathcal{R}_{AT}}{\partial \rho^2} \left(1 - \frac{1}{\sqrt{2}}, -1 + \frac{1}{\sqrt{2}}\right) \frac{\partial^2 \mathcal{R}_{AT}}{\partial \beta^2} \left(1 - \frac{1}{\sqrt{2}}, -1 + \frac{1}{\sqrt{2}}\right) - \left[\frac{\partial^2 \mathcal{R}_{AT}}{\partial \rho \beta} \left(1 - \frac{1}{\sqrt{2}}, -1 + \frac{1}{\sqrt{2}}\right)\right]^2 = \frac{(32)^3}{(81)^2} > 0. \tag{190}$$

Since $\frac{\partial^2 \mathcal{R}_{AT}}{\partial \beta^2} \left(1 - \frac{1}{\sqrt{2}}, -1 + \frac{1}{\sqrt{2}}\right) = -\frac{1024}{87} < 0$, it holds that $\left(1 - \frac{1}{\sqrt{2}}, -1 + \frac{1}{\sqrt{2}}\right)$ is a relative minima for \mathcal{R}_{AT} . Similarly, $\left(1 + \frac{1}{\sqrt{2}}, -1 - \frac{1}{\sqrt{2}}\right)$ is a relative minima for \mathcal{R}_{AT} .

As for G_{11} , we have $\frac{\partial G_{11}}{\partial \rho} = \frac{(2(1-\rho)^2+1)(1-\rho)(1+\beta)}{(1-\rho)^2(1+(1-\rho)^2)^{\frac{5}{2}}(1+(1+\beta)^2)^{\frac{3}{2}}}$, $\frac{\partial G_{11}}{\partial \beta} = \frac{(1-2(1+\beta)^2)(1-\rho)(1+\beta)}{(1-\rho)(1+(1-\rho)^2)^{\frac{3}{2}}(1+(1+\beta)^2)^{\frac{5}{2}}}$ *Eq. (167)* and *Eq. (168)*. Hence, the critical point are $(\rho_{critical}, \beta_{critical}) = \left(1 - \frac{1}{\sqrt{2}}, 0\right), \left(1 + \frac{1}{\sqrt{2}}, -2\right)$. Clearly, $\beta_{critical} = 0, -2$ are never permissible since $M/G/1$ is stable. Thus, no conclusion can be drawn for G_{11} .

Engaging the same procedure, it could be verified that both maxima and minima for all the remaining components of the Spacetime curvature (Einstein Tensor) subject to the Angular Technique, \wp_{AT} is undecidable.

Theorem 16. The underlying QM satisfies:

- I. Both maxima and minima of \mathcal{R}_{MT} is undecidable.
- II. Both maxima and minima for all the components of the Spacetime curvature (Einstein Tensor) subject to the Monge Technique, \wp_{MT} is undecidable.

Proof:

I. $\mathcal{R}_{MT} = -2$ *Eq. (171)*. Hence, $\frac{\partial \mathcal{R}_{AT}}{\partial \rho} = 0 = \frac{\partial \mathcal{R}_{AT}}{\partial \beta}$ for all ρ, β . It can be shown that D of *Eq. (45)* is zero. Hence, I) follows.

II. The proof is immediate for G_{12} , since it is zero *Eq. (175)*. By *Eq. (174)*, $G_{11} = \frac{(1-\alpha)}{(1-\rho)^2} = G_{21}$. We have $\frac{\partial G_{11}}{\partial \rho} = \frac{2(1-\alpha)}{(1-\rho)^3}$, $\frac{\partial G_{11}}{\partial \alpha} = \frac{-1}{(1-\rho)^2}$ (=0 if and only if $\rho \rightarrow \infty$). Hence, both maxima and minima are undecidable for G_{11}, G_{21} . Finally, $G_{22} = -\frac{1}{(1+\beta)^2}$ *Eq. (176)*. $\frac{\partial G_{22}}{\partial \beta} = \frac{2}{(1+\beta)^3}$. Therefore, $\frac{\partial G_{22}}{\partial \beta} = 0$ if and only if $\beta \rightarrow \infty$. Consequently, it is not possible to decide maxima and minima for G_{22} .

The following theorem captures the impact of stability(instability) of M/G/1 QM on the increasability (decreasability) of the only non-zero component of RCT, $R_{21}^{(0)}$.

Theorem 17. The underlying QM satisfies:

- I. $R_{21}^{(0)}$ is forever increasing in $\rho \Leftrightarrow$ the underlying QM is unstable.
- II. $R_{21}^{(0)}$ forever decreases in ρ if and only if M/G/1 QM is stable.

Proof:

I. Necessity:

Assume that M/G/1 QM is unstable, then $\rho > 1$. We have $\frac{\partial}{\partial \rho}(R_{21}^{(0)}) = -\frac{2}{(1-\rho)^3}$ (8.48). By $\rho > 1$, $\frac{\partial}{\partial \rho}(R_{21}^{(0)}) > 0$, which directly implies by the Preliminary Theorem, that $R_{21}^{(0)}$ is forever increasing in ρ .

Sufficiency

Let $R_{21}^{(0)}$ be forever increasing in ρ , then by the Preliminary Theorem, $\frac{\partial}{\partial \rho}(R_{21}^{(0)}) = -\frac{2}{(1-\rho)^3} > 0$. This implies $(\rho - 1)^3 > 0$. Hence, $\rho > 1$. This proves I).

Engaging the same procedure, II) follows.

The following Theorem captures the dual impact between (IG) and Queueing Theory. This dual impact is influencing the existence of FIM, upon the behaviour of $\Delta_{[g^{ij}]}$.

Theorem 18. The underlying QM satisfies:

- I. $\Delta_{[g^{ij}]}$ is forever increasing(decreasing) in $\rho \Leftrightarrow$ the underlying QM is (stable) unstable.
- II. $\Delta_{[g^{ij}]}$ is forever increasing(decreasing) in $\beta \Leftrightarrow$ the underlying QM is (unstable) stable.
- III. The inflection point of $\Delta_{[g^{ij}]}$ with respect to β is at $\rho = 1$, where $\Delta_{[g^{ij}]}$ changes its behaviour around this threshold $\rho = 1$.

Proof:

$$1) \text{ We have } [g^{ij}] = \frac{\text{adj}[g_{ij}]}{\Delta} = \begin{pmatrix} (1-\rho)^2 & 0 \\ 0 & -(\beta+1)^2 \end{pmatrix} \text{ Theorem 3. Therefore,} \\ \Delta_{[g^{ij}]} = -(\beta+1)^2(1-\rho)^2. \quad (191)$$

By Eq. (191), $\Delta_{[g^{ij}]} = 0$ if and only if $\rho = 0$ or $\beta = -1$ (this is never permissible).

$$\frac{\partial \Delta_{[g^{ij}]}}{\partial \rho} = 2(1-\rho)(\beta+1)^2. \quad (192)$$

Clearly, from Eq. (192), it follows that:

$$\frac{\partial \Delta_{[g^{ij}]}}{\partial \rho} > 0 (< 0) \text{ if and only if } (\rho < 1) (\rho > 1). \quad (193)$$

By PT 2.15, the proof of 1) follows.

2) We have

$$\frac{\partial \Delta_{[g^{ij}]}}{\partial \beta} = -2(1-\rho)^2(\beta+1), \quad (194)$$

$$\frac{\partial \Delta_{[g^{ij}]}}{\partial \beta} > 0 (< 0) \text{ if and only if } (\rho > 1) (\rho < 1). \quad (195)$$

By (PT) 2.15, the proof of 2) follows.

3) It is straightforward to see that

$$\frac{\partial^2(\Delta_{[g^{ij}]}(\rho, \beta))}{\partial \beta^2} = -2(1 - \rho)^2. \tag{196}$$

$\frac{\partial^2(\Delta_{[g^{ij}]}(\rho, \beta))}{\partial \beta^2}$ if and only if $\rho = 1$, which proves the required statement.

DATA $\Delta_{[g^{ij}]}$; $\Delta_{[g^{ij}]}$ for $\beta = 2, \rho \in (0,1)$.

Case 1.




Fig. 23. Impact of stability of M/G/1 queue on (-det FIM), for SCV =2.

As observed in Fig. 23, det (IFIM) is increasing in server utilization if and only if M/G/1 QM is stable. Also, this proves how the stability of the underlying QM impacts the existence of $[g^{ij}]$.

Case 2. $\Delta_{[g^{ij}]}$ for $\beta = 2, \rho \in [1, \infty)$.




Fig. 24. Impact of instability of M/G/1 queue on (-det FIM), for SCV =2.

As observed in Fig. 24, instability of M/G/1 QM is unstable $\Leftrightarrow \Delta$ decreases in ρ .

Case 3. $\Delta_{[g^{ij}]}$ behaviour for β in stability phase of M/G/1 QM, $\rho = 0.5$.




Fig. 25. Impact of stability of M/G/1 queue on $(-\det FIM)$, for $SU = 0.5$.

As seen above in Fig. 25, $\det (IFIM)$ is forever decreasing in SCV if and only if M/G/1 QM is stable.

Case 4. $\Delta_{[g^{ij}]}$ behaviour for β in instability phase of MG1 QM, $\rho = 2$




Fig. 26. Impact of instability of M/G/1 queue on $(-\det FIM)$, for $SU = 0.5$.

As observed in Fig. 26, within the instability phase of M/G/1 QM, $\beta \in (0,1)$, $\det (IFIM)$ is decreasing in β .

Case 5.




Fig. 27. Impact of SU on the second derivative of (det FIM) with respect to SCV.

Fig. 27 justifies that the threshold of stability of M/G/1 QM, $\rho = 1$, would be the inflection point of det (IFIM) as well as being the decision parameter that controls the existence of inverse of (IFIM).

Define (QIGUs) by the triad functions, namely $h_1^{QIGU}, h_2^{QIGU}, h_3^{QIGU}$, with $h_1^{QIGU}, h_2^{QIGU}: M/G/1 QM \rightarrow [g_{ij}]$, where $[g_{ij}] = \begin{pmatrix} \frac{1}{(1-\rho)^2} & 0 \\ 0 & \frac{-1}{(\beta+1)^2} \end{pmatrix}$ Theorem 3.

$$h_1^{QIGU}(\rho) = g_{11} = \frac{1}{(1-\rho)^2}, \rho \in (0,1), \tag{197}$$

$$h_2^{QIGU}(\beta) = -g_{22} = \frac{1}{(1+\beta)^2}, \beta \in (1, \infty), \tag{198}$$

$$h_3^{QIGU}(\rho, \beta) = \begin{cases} \Psi_{n=0}(\rho, \beta) = -\ln(1 - \rho), & n = 0, \\ \Psi_{n>0}(\rho, \beta) = \ln(\beta + 1) - \ln(1 - \rho) - \ln 2, & n > 0. \end{cases} \tag{199}$$

provided that $\Psi_{n=0}(\rho, \beta), \Psi_{n>0}(\rho, \beta)$ are determined by Eq. (54), Eq. (61) of Theorem 3 respectively.

9 | Queueing Theoretic Impact on the Continuity of New Devised Queueing-Information Geometric Unifiers (QIGU)

Theorem 19. For the stable M/G/1 QM

- I. h_1^{QIGU} is continuous, for $\rho \in (0,1)$.
- II. h_1^{QIGU} is well-defined.
- III. h_1^{QIGU} is one-to-one.
- IV. h_1^{QIGU} is onto.
- V. The inverse function of h_1^{QIGU} is characterized by, $(h_1^{QIGU})^{-1}(\rho) = 1 - \frac{1}{\sqrt{\rho}}, \rho \in (0,1)$.
- VI. h_2^{QIGU} is continuous, for $\beta \in (1, \infty)$.
- VII. h_2^{QIGU} is well-defined.

VIII. h_2^{QIGU} is one-to-one.

IX. h_2^{QIGU} is onto.

X. The inverse function of h_2^{QIGU} is characterized by, $(h_2^{\text{QIGU}})^{-1}(\beta) = -1 + \frac{1}{\sqrt{\beta}}$, $\beta \in (1, \infty)$.

XI. $\Psi_{n=0}(\rho, \beta)$ is well-defined.

XII. $\Psi_{n=0}(\rho, \beta)$ is continuous if and only if M/G/1 QM is stable.

XIII. $\Psi_{n=0}(\rho, \beta)$ is discontinuous \Leftrightarrow the instability of the underlying QM is satisfied.

XIV. $\Psi_{n=0}(\rho, \beta)$ is one-to-one.

XV. $\Psi_{n=0}(\rho, \beta)$ is onto.

XVI. The inverse function of $\Psi_{n=0}(\rho, \beta)$ is characterized by, $(\Psi_{n=0}(\rho, \beta))^{-1}(\rho) = 1 - e^{-\rho}$, $\rho \in (0, 1)$.

XVII. $\Psi_{n>0}(\rho, \beta)$ characterizes a family of non-well-defined (ρ, β) dependent functions.

XVIII. $\Psi_{n>0}(\rho, \beta)$ is continuous if and only if M/G/1 QM is stable.

XIX. $\Psi_{n>0}(\rho, \beta)$ is discontinuous \Leftrightarrow the instability of the underlying QM is satisfied.

Proof

1) Let $\rho_1, \rho_2 \in (0, 1)$ such that there is a $\delta > 0$ satisfying

$$|\rho_1 - \rho_2| < \delta, \quad (200)$$

$$|h_1^{\text{QIGU}}(\rho_1) - h_1^{\text{QIGU}}(\rho_2)| = \left| \frac{1}{(1-\rho_1)^2} - \frac{1}{(1-\rho_2)^2} \right| = \left| \frac{(\rho_1 - \rho_2)(\rho_1 + \rho_2 - 2)}{(1-\rho_1)^2(1-\rho_2)^2} \right| < \frac{\delta|\rho_1 + \rho_2 - 2|}{(1-\rho_1)^2(1-\rho_2)^2} = \varepsilon. \quad (201)$$

Clearly, by Eq. (46), proof of I follows.

It is also clear that if either $\rho_1 = 1$ or $\rho_2 = 1$. Then, the underlying QM is unstable, which implies by Eq. (201) that $\varepsilon \rightarrow \infty$. This implies that h_1^{QIGU} is discontinuous at $\rho = 1$.

II. Let $\rho_1, \rho_2 \in (0, 1)$, $\rho_1 \neq \rho_2$. Assume that $h_1^{\text{QIGU}}(\rho_1) = h_1^{\text{QIGU}}(\rho_2)$. Hence, $\frac{1}{(1-\rho_1)^2} = \frac{1}{(1-\rho_2)^2}$, equivalently, $(1 - \rho_1)^2 = (1 - \rho_2)^2$. This means, $\rho_1 = \rho_2$ or $\rho_1 + \rho_2 = 2$. We have to reject, $\rho_1 = \rho_2$. Also, $\rho_1 + \rho_2 = 2$ is impossible since $\rho_1, \rho_2 \in (0, 1)$, Clearly, a contradiction follows. Therefore, II holds.

III. $h_1^{\text{QIGU}}(\rho_1) = h_1^{\text{QIGU}}(\rho_2)$, this implies that $\rho_1 = \rho_2$.

IV. Obviously, every $\rho \in (0, 1)$ uniquely characterizes $\frac{1}{(1-\rho)^2}$ such that $h_1^{\text{QIGU}}(\rho) = g_{11} = \frac{1}{(1-\rho)^2}$. Hence, h_1^{QIGU} is onto.

V. Assume $h_1^{\text{QIGU}}(\rho) = y$. Hence,

$$(h_1^{\text{QIGU}})^{-1}(\rho) = 1 - \frac{1}{\sqrt{\rho}}, \rho \in (0, 1). \quad (202)$$

VI. Let $\beta_1, \beta_2 \in (1, \infty)$ such that there is a $\delta > 0$ satisfying:

$$|\beta_1 - \beta_2| < \delta, \quad (203)$$

$$|h_2^{\text{QIGU}}(\beta_1) - h_2^{\text{QIGU}}(\beta_2)| = \left| \frac{1}{(1+\beta_1)^2} - \frac{1}{(1+\beta_2)^2} \right| = \left| \frac{(\beta_1 - \beta_2)(\beta_1 + \beta_2 - 2)}{(1+\beta_1)^2(1+\beta_2)^2} \right| < \frac{\delta}{16} |\beta_1 + \beta_2 - 2| = \varepsilon. \quad (204)$$

Clearly by Eq. (46), the proof of VI follows.

VII. Let $\beta_1, \beta_2 \in (1, \infty)$, $\beta_1 \neq \beta_2$. Assume that $h_2^{\text{QIGU}}(\beta_1) = h_2^{\text{QIGU}}(\beta_2)$. Hence, $\frac{1}{(1+\beta_1)^2} = \frac{1}{(1+\beta_2)^2}$, equivalently, $(1 + \beta_1)^2 = (1 + \beta_2)^2$. This means, $\beta_1 = \beta_2$ or $\beta_1 + \beta_2 = -2$. We must reject, $\beta_1 = \beta_2$. Also, $\beta_1 + \beta_2 = -2$ is impossible since $\beta_1, \beta_2 \in (1, \infty)$, Clearly, a contradiction follows. Therefore, holds.

It is also clear that even if the underlying QM is unstable (or equivalently, $\beta \in (0,1]$, which implies by Eq. (104), that $\varepsilon \rightarrow \infty$). This implies that h_2^{QIGU} is everywhere continuous.

VIII. Assume $h_2^{\text{QIGU}}(\beta) = \frac{1}{(1+\beta)^2} = w, \beta \in (1, \infty)$. This leaves one choice,

$$\beta = -1 + \frac{1}{\sqrt{w}}. \quad (205)$$

This shows that, $(h_2^{\text{QIGU}})^{-1}(\beta) = 1 - \frac{1}{\sqrt{\beta}}, \beta \in (1, \infty)$.

X. Let $\rho_1, \rho_2 \in (0,1), \rho_1 \neq \rho_2$. Assume that $\Psi_{n=0}(\rho_1) = \Psi_{n=0}(\rho_2)$. Hence, $-\ln(1 - \rho_1) = -\ln(1 - \rho_2)$, equivalently, $(1 - \rho_1) = (1 - \rho_2)$. This means, $\rho_1 = \rho_2$. Clearly, a contradiction follows. Therefore, X holds.

XI. Necessity: Assume that M/G/1 QM is stable, then $\rho_1, \rho_2 \in (0,1)$.

Suppose that there is a $\delta > 0$ satisfying:

$$|\rho_1 - \rho_2| < \delta, \quad (206)$$

$$|\Psi_{n=0}(\rho_1) - \Psi_{n=0}(\rho_2)| = |\ln(1 - \rho_1) - \ln(1 - \rho_2)|. \quad (207)$$

It is well known that Maclaren's series of $\ln(1 - \rho)$, for $\rho \in (0,1)$ is determined by

$$\ln(1 - \rho) = -\sum_{n=1}^{\infty} \frac{\rho^n}{n}. \quad (47)$$

Thus, it follows that:

$$|\ln(1 - \rho_1) - \ln(1 - \rho_2)| < \delta \sum_{n=1}^{\infty} (\rho_1^{n-1} + \rho_1^{n-2} + \dots + 1) = \frac{\delta}{(1-\rho_1)} = \varepsilon. \quad (208)$$

This proves continuity.

Sufficiency:

By Eq. (208), there exists $\frac{\delta_1}{(1-\rho_1)} = \varepsilon_1 > 0, \frac{\delta_2}{(1-\rho_2)} = \varepsilon_2 > 0$, with $\delta = \min(\delta_1, \delta_2) > 0, \varepsilon = \min(\varepsilon_1, \varepsilon_2)$ satisfying, Eq. (206) and Eq. (208). Consequently, $\rho_1, \rho_2 \in (0,1)$, which directly implies the underlying QM's stability.

XII. Following Eq. (208), discontinuity of $\Psi_{n=0}$ occurs if and if $\rho_1 \geq 1$ to enforce ε to be infinite of any negative real number. Therefore, XII holds.

The validity of both XIII and XIV can easily be verified.

Assume $\Psi_{n=0}(\rho) = -\ln(1 - \rho) = m, \rho \in (0,1)$. Thus, it holds that $1 - \rho = e^{-m}$. Consequently, $\rho = 1 - e^{-m}$. This shows that $(\Psi_{n=0})^{-1}(\rho) = 1 - e^{-\rho}, \rho \in (0,1)$, which proves XV.

XVI. Let $\rho_1, \rho_2 \in (0,1), (\beta_1, \beta_2) \in (1, \infty)$ such that $\rho_1 \neq \rho_2, \beta_1 \neq \beta_2$. Assume that $\Psi_{n>0}(\rho_1, \beta_1) = \Psi_{n>0}(\rho_2, \beta_2)$. Hence,

$$\frac{(\beta_1+1)}{(1-\rho_1)} = \frac{(\beta_2+1)}{(1-\rho_2)}. \text{ This means,}$$

$$(\beta_1 - \beta_2) + (\rho_1 - \rho_2) = (\beta_2\rho_1 - \beta_1\rho_2). \quad (209)$$

By Eq. (209), XVI) holds.

It is obvious that $\Psi_{n>0}$ is continuous for all $\beta \in (1, \infty), \rho \in (0,1)$. This proves the necessity requirement. As for the sufficiency requirement, let $\Psi_{n>0}(\rho, \beta)$ be continuous. This directly implies that $\ln(1 - \rho)$ attains non-infinite real values. Consequently, $1 - \rho > 0$. Hence, the underlying M/G/1 QM is stable. This proves 18).

To prove XIX, it is clear that $\Psi_{n>0}(\rho, \beta)$ is discontinuous if and only if both $\ln(\beta + 1)$ and $\ln(1 - \rho)$ is discontinuous. This is equivalent to $\beta > -1, \rho \geq 1$. Equivalently, M/G/1 QM is unstable.

10 | Closing Remarks with the Next Phase of Research

This paper discusses the application of information geometric concepts in queueing theory, specifically focusing on the specified QM. It introduces the Fisher information metric, the α -connection, and analyses the geometric properties of the queue manifold, such as Gaussian and Ricci curvatures. The paper also highlights the potential for further research in IG unification with existing knowledge of scientific fields.

References

- [1] Mageed, I. A., Zhou, Y., Liu, Y., & Zhang, Q. (2023). Towards a revolutionary info-geometric control theory with potential applications of fokker planck kolmogorov (FPK) equation to system control, modelling and simulation. *2023 28th international conference on automation and computing (ICAC)* (pp. 1-6). IEEE. <https://doi.org/10.1109/ICAC57885.2023.10275271>
- [2] Mageed, I. A., Yin, X., Liu, Y., & Zhang, Q. (2023). $\mathbb{Z}_{[a, b]}$ of the stable five-dimensional M/G/1 queue manifold formalism's info-geometric structure with potential info-geometric applications to human computer collaborations and digital twins. *2023 28th International conference on automation and computing (ICAC)* (pp. 1-6). IEEE. DOI: 10.1109/ICAC57885.2023.10275185
- [3] Mageed, I. A. (2024). *Info-geometric analysis of gamma distribution manifold with gamma distribution impact to advance satellite earth observations*. <https://www.preprints.org/manuscript/202402.0968>
- [4] Mageed, I. A., Zhang, Q., Akinci, T. C., Yilmaz, M., & Sidhu, M. S. (2022). Towards abel prize: the generalized brownian motion manifold's fisher information matrix with info-geometric applications to energy works. *2022 global energy conference (GEC)* (pp. 379-384). IEEE. <https://doi.org/10.1109/GEC55014.2022.9987239>
- [5] Mageed, I. A. (2024). Info-geometric analysis of the stable queue manifold dynamics with queue applications to e-health. <https://www.preprints.org/manuscript/202401.1813>
- [6] Mageed, I. A., & Kouvatso, D. D. (2021). The impact of information geometry on the analysis of the stable M/G/1 queue manifold. *ICORES* (pp. 153-160). DOI: 10.5220/0010206801530160
- [7] Mageed, I. A., & Kouvatso, D. D. (2019). Information geometric structure of stable m/g/1 queue manifold and its matrix exponential. *35th UK performance engineering workshop 16 December 2019* (p. 116). https://www.researchgate.net/profile/Dr-Vijay-Singh-2/publication/340951179_proceedings/links/5ea70fb145851553fab34b07/proceedings.pdf#page=123
- [8] Miyagawa, S. (2010). *MIT open course ware: How it began*. <https://www.nae.edu/162657/MIT-OpenCourseWare-How-it-Began>
- [9] Mageed, I. A. (2024). *On the Rényi entropy functional, tsallis distributions and lévy stable distributions*. https://www.preprints.org/manuscript/202401.1996/download/final_file
- [10] Grivel, E., Diversi, R., & Merchan, F. (2021). Kullback-Leibler and Rényi divergence rate for Gaussian stationary ARMA processes comparison. *Digital signal processing*, 116, 103089.
- [11] Cai, T. T., Guo, Z., & Ma, R. (2023). Statistical inference for high-dimensional generalized linear models with binary outcomes. *Journal of the american statistical association*, 118(542), 1319–1332. <https://doi.org/10.1080/01621459.2021.1990769>
- [12] Marmin, A., de Morais Goulart, J. H., & Févotte, C. (2023). Majorization-minimization for sparse nonnegative matrix factorization with the β -divergence. *IEEE transactions on signal processing*, 71, 1435-1447. <https://arxiv.org/pdf/2207.06316>
- [13] Cilingir, H. K., Manzelli, R., & Kulis, B. (2020). Deep divergence learning. *International conference on machine learning* (pp. 2027-2037). PMLR. <http://proceedings.mlr.press/v119/cilingir20a/cilingir20a.pdf>
- [14] Chow, B., Lu, P., & Ni, L. (2023). *Hamilton's Ricci flow* (Vol. 77). American mathematical society, Science Press. <https://books.google.com/books>
- [15] Gilkey, P., Brozos-Vázquez, M., Garcia-Río, E., Nikčević, S., & Vásquez-Lorenzo, R. (2022). *The geometry of Walker manifolds*. Springer Nature.

- [16] Bachmann, G., Bécigneul, G., & Ganea, O. (2020). Constant curvature graph convolutional networks. In *International conference on machine learning* (pp. 486-496). PMLR. <http://proceedings.mlr.press/v119/bachmann20a/bachmann20a.pdf>
- [17] Sun, A. (2020). Compactness of constant mean curvature surfaces in a three-manifold with positive Ricci curvature. *Pacific journal of mathematics*, 305(2), 735–756. <https://arxiv.org/pdf/1804.09328>
- [18] Meyer, C. D. (2023). Matrix analysis and applied linear algebra. SIAM. <https://epubs.siam.org/doi/pdf/10.1137/1.9781611977448.bm>
- [19] Zhang, Y., & Zheng, J. (2022). An overview of developable surfaces in geometric modeling. *Recent patents on engineering*, 16(5), 87–103. <https://doi.org/10.2174/1872212115666210512021156>
- [20] Kamarudzaman, A. S. M., Nasir, N. H. M., & Misro, M. Y. (2022). Gaussian and mean curvature of biquintic trigonometric bézier surface. *Pertanika journal of science & technology*, 30(2). https://www.researchgate.net/profile/Md-Yushalify-Misro/publication/359897709_Gaussian_and_Mean_Curvature_of_Biquintic_Trigonometric_Bezier_Surface/links/6273d691b1ad9f66c8a41971/Gaussian-and-Mean-Curvature-of-Biquintic-Trigonometric-Bezier-Surface.pdf
- [21] Wang, J., Xing, H., Jin, Z., Tan, Y., Pang, S., & Liu, J. (2023). 3D geometric modelling of the northwest pacific slabs: a review and new high-precision model. *Earth-science reviews*, 238, 104351. <https://doi.org/10.1016/j.earscirev.2023.104351>
- [22] Banica, T. (2024). Calculus and applications. ArXiv preprint arxiv:2401.00911. <https://arxiv.org/pdf/2401.00911>
- [23] Latayada, M. J. R. (2023). On the number of restricted one-to-one and onto functors having integral coordinates. *European journal of pure and applied mathematics*, 16(4), 2751–2762. <http://www.ejpam.com/index.php/ejpam/article/view/4901/1532>
- [24] Irving, R. S. (2013). Beyond the quadratic formula (Vol. 43). MAA. <https://books.google.com/books>
- [25] Mageed, I. A., & Zhang, Q. (2023). Threshold theorems for the tsallisian and rényian (tr) cumulative distribution functions (CDFs) of the heavy-tailed stable $m/g/1$ queue with tsallisian and rényian entropic applications to satellite images (SIs). *Electronic journal of computer science and information technology*, 9(1), 41–47. <https://doi.org/10.52650/ejcsit.v9i1.161>
- [26] Roussel, M. R. (2005). Stability analysis for ODEs. Nonlinear dynamics, lecture notes, university hall, canada, 78. <https://fac.ksu.edu.sa/sites/default/files/stability.pdf>
- [27] Schloss, J., Barnett, P., Sachdeva, R., & Busch, T. (2020). Controlled creation of three-dimensional vortex structures in Bose-Einstein condensates using artificial magnetic fields. *Physical review a*, 102(4), 43325. <https://doi.org/10.1103/PhysRevA.102.043325>
- [28] Bailes, M., Berger, B. K., Brady, P. R., Branchesi, M., Danzmann, K., Evans, M., ...& others. (2021). Gravitational-wave physics and astronomy in the 2020s and 2030s. *Nature reviews physics*, 3(5), 344–366. <https://doi.org/10.1038/s42254-021-00303-8>
- [29] Dawkins, P. (2021). *Paul's Online Notes, web source*. National council of teachers of mathematics. <https://tutorial.math.lamar.edu/>
- [30] Breneman, J. E., Sahay, C., & Lewis, E. E. (2022). *Introduction to reliability engineering*. John Wiley & Sons.
- [31] Mageed, I. A., & Zhang, Q. (2022). An information theoretic unified global theory for a stable $m/g/1$ queue with potential maximum entropy applications to energy works. *2022 global energy conference (GEC)* (pp. 300-305). IEEE. <https://doi.org/10.1109/GEC55014.2022.9986719>